

“FACTORS RELATED TO FALLS AMONG THE ELDERLY IN THE EASTERN THAILAND”

Ploi Kongkoot
Thailand

56th Master of Public Health/International Course in Health Development

*KIT (Royal Tropical Institute)
Vrije Universiteit Amsterdam (VU)*

"FACTORS RELATED TO FALLS AMONG THE ELDERLY IN THE EASTERN THAILAND"

A thesis submitted in partial fulfilment of the requirement for the degree of
Master of Science in Public Health

by
Ploi Kongkoot

Thailand

Declaration:

Where other people's work has been used (from either a printed source, internet or any other source), this has been carefully acknowledged and referenced in accordance with departmental requirements.

The thesis "FACTORS RELATED TO FALLS AMONG THE ELDERLY IN THE EASTERN THAILAND" is my own work.

Signature:



56th Master of Public Health/International Course in Health Development
(MPH/ICHD)

16 September 2019 – 4 September 2020

KIT (Royal Tropical Institute)/Vrije Universiteit Amsterdam

Amsterdam, The Netherlands September 2020

Organised by:

KIT (Royal Tropical Institute) Amsterdam, The Netherlands

In co-operation with:

Vrije Universiteit Amsterdam (VU) Amsterdam, The Netherlands

Table of Contents

<i>Lists of tables and figures</i>	<i>IV</i>
Table.....	<i>IV</i>
Figure.....	<i>IV</i>
<i>Abbreviations</i>	<i>V</i>
<i>Key Terms</i>	<i>V</i>
<i>Abstract</i>	<i>VI</i>
<i>Introduction</i>	<i>VII</i>
<i>CHAPTER ONE : BACKGROUND OF THE COUNTRY</i>	<i>1</i>
<i>CHAPTER TWO : PROBLEM STATEMENT AND METHODOLOGY</i>	<i>4</i>
2.1 Problem statement	<i>4</i>
2.2 Justification	<i>5</i>
2.3 Research objective	<i>6</i>
2.3.1 General objective.....	<i>6</i>
2.3.2 Specific objective	<i>6</i>
2.4 Methodology and analytical framework.....	<i>6</i>
2.4.1 Secondary data analysis	<i>6</i>
2.4.2 Literature review	<i>8</i>
2.4.3 Analytical Framework.....	<i>9</i>
<i>CHAPTER THREE : STUDY FINDING</i>	<i>11</i>
3.1 Finding introduction.....	<i>11</i>
3.2 Description of the characteristics of respondents.....	<i>11</i>
3.3 Individual cause	<i>12</i>
3.3.1Dependence factors.....	<i>12</i>
3.3.2 Independence factors	<i>19</i>
3.4 Proximal cause	<i>22</i>
3.4.1 Environmental factor	<i>22</i>
3.4.2 Family factors	<i>26</i>
3.5 Distal Cause.....	<i>28</i>
3.5.1 Social and community support.....	<i>28</i>
3.5.2 Health system, Policy and Government factors.....	<i>28</i>
<i>CHAPTER FOUR : FALLS PREVENTION AND BEST PRACTICES</i>	<i>31</i>
4.1 Prevention at the individual level	<i>31</i>
4.1.1 Motor and balance problems.....	<i>31</i>
4.1.2 Correct visual impairment by Cataract surgery	<i>31</i>
4.2 Prevention at the proximal level.....	<i>31</i>
4.3 Prevention at the distal level	<i>32</i>
<i>CHAPTER FIVE : DISCUSSION AND LIMITATION</i>	<i>33</i>
5.1 Discussion	<i>33</i>
5.1.1 General.....	<i>33</i>

5.1.2 Individual level	33
5.1.3 Proximal level	34
5.1.4 Distal level	35
5.1.5 Preventive interventions of falls	36
5.2 Limitations of the study	36
CHAPTER SIX : CONCLUSION AND RECOMMENDATION	38
6.1 Conclusion.....	38
6.2 Recommendation.....	39
References:.....	40
Acknowledgement	45
Annex	46
Annex 1.....	46
Annex 2.....	46
Annex 3.....	47
Annex 4.....	48
Annex 5.....	50
Annex 6.....	51

Lists of tables and figures

Table

Table 1 Demographic data of study population among 450 respondents age >60 years old in a survey in Eastern Thailand in 2018	11
Table 2 Descriptive data of education, marital status, religion, occupation and financial status among total 450 participants age >60 in the survey in Eastern of Thailand in 2018...	13
Table 3 Bivariate analysis of dependent individual factors with falls in a survey in Eastern Thailand 2018	14
Table 4 Descriptive data of Physical and health status among total 450 participants age >60 in the survey in Eastern of Thailand in 2018	16
Table 5 Descriptive data of medical conditions among total 450 participants age >60 in the survey in Eastern of Thailand in 2018.....	17
Table 6 Bivariate analysis of dependent individual factors with falls in a survey in Eastern Thailand 2018	18
Table 7 Descriptive data of healthy life style among total 450 participants age >60 in the survey in Eastern of Thailand in 2018.....	19
Table 8 Bivariate analysis of independent individual factors with falls in a survey in Eastern Thailand in 2018	21
Table 9 Descriptive data of living condition and area around houses of total 450 participants age >60 in the survey in Eastern of Thailand in 2018.....	22
Table 10 Bivariate analysis of indoor environmental factors with falls in a survey in Eastern Thailand 2018	23
Table 11 Bivariate analysis of outdoor environmental factors with falls in a survey in Eastern Thailand 2018	25
Table 12 Descriptive data of living condition and family support of total 450 participants age >60 in the survey in Eastern of Thailand in 2018	26
Table 13 Bivariate analysis of proximal factors with falls in a survey in Eastern Thailand 2018	27
Table 14 Descriptive data of social and community support of total 450 participants age >60 in the survey in Eastern of Thailand in 2018.....	28
Table 15 Descriptive data the health insurance and support from government of total 450 participants age >60 in the survey in Eastern of Thailand in 2018	28
Table 16 Bivariate analysis of distal factors with falls in a survey in Eastern Thailand 2018 .	29

Figure

Figure 1 The map of Eastern Thailand, Figure 2 Provinces in Eastern Thailand	1
Figure 3 Change in Crude Birth Rate, Mortality Rate and Natural Growth in Thailand 2018...	2
Figure 4 Young Children and Older People as a percentage of Global population,1950-5050	4
Figure 5 The Mortality rate of falls Among the elderly and other age groups of the population from 2007-2016 in Thailand	5
Figure 6 Causation of falls conceptual framework	10
Figure 7 Causes of falls pie chart, among 121 respondents age >60 years old who had fallen in the past year in a survey in Eastern Thailand 2018	12

Abbreviations

BMI	Body Mass Index
EEC	Eastern Economic Corridor
NSO	National Statistic Office
NCD	Non-Communicable Disease
GDP	Gross Domestic Product
WHO	World Health Organization
VU	Vrije Universiteit Amsterdam (University in Amsterdam)
NHS	National Health Service
MeSH	Medical Subject HEading
HF	Health facility
SD	Standard deviation
OR	Odd ratio
DM	Diabetes Mellitus
HT	Hypertension
IHD	Ischemic heart disease
DLP	Dyslipidemia
Knee OA	Knee osteoarthritis
HL	Health literacy
ESA	Elderly subsistence allowance
AOR	Adjusted Odds Ratio
IRR	Incidence Rate Ratio
RaR	Rate Ratio

Key Terms

Pes planus	Known as flatfoot, which is the deformity of the foot when it abnormally flattened.
Plantar fasciitis	Inflammation at the ligament on the bottom of the heel.
Parkinson's disease	"A long-term generative disorder of the central nervous system and mainly affect to movement function".
Dementia	"A syndrome that affect to deterioration in memory, thinking, behaviour and the ability to perform everyday activities. Usually Affect old age group."
Health literacy	"The achievement of a level of knowledge, personal skills and confidence to take action to improve health by changing personal lifestyles and living conditions."

Abstract

Title: Factors related to falls among the elderly in Eastern Thailand

Background: Falls, have a high incidence, are a major health problem among the elderly due to the ageing society change in Thailand.

Method: A mixed study of Secondary data analysis and literature review was used to find the factors related to the falls. The data from the survey among 450 elders age 60 and above, who lived in Eastern Thailand in 2018, will be analyzed between fallers and non-fallers.

Result: Participants in the survey of the age 60-88 years old were 450 elders, from that 121 participants had fallen within 1 year (26.9%), these are 19 men and 102 women. Among fallers, participants with an age more than 80 had the highest proportion of falls. In individual level the reported BMI ($p=0.038$), excessive waist circumference ($p<0.001$), hearing impairment ($p=0.037$), balance impairment ($p<0.001$), and taking sleeping pills ($p=0.027$) were significantly associated with falls. In proximal level it was found that the area around houses, with rough ground and rocks, clear areas, and with incline pathways, were associated to falls. There is no report of association in distal level. Exercise, home hazardous modification and combined interventions were found effective to prevent falls among the elderly.

Conclusion: Risks factor of falls were found because of different levels of causes. Physical status in the individual level are related to falls as environmental hazardous in the proximal level. Preventive interventions, exercise and home hazard modification, are mandatory to decrease fall incidences in elders.

Keyword: falls, elderly, prevention, risk factors, Eastern Thailand

Word count :12498

Introduction

I am a medical doctor in Health Center 6, Thailand, with six-year experience in the clinical field. My work experience gave me not only medical skills, but I have earned communication and research skills. In the clinical field, working experiences brought me the idea of expanding my skills and knowledge in health promotion, prevention diseases, and research on health issues. I found these are crucial in the health system as same as the medical treatment and I always believe in "Prevention is better than cure." My workplace belongs to The Department of Health, which responsibilities are providing medical treatment, doing health research and monitoring and evaluating health care indicators in a specific area. I had contributed to research, projects, and a few were done in the ageing issue. Nowadays, Thai MoPH has concerned ageing health problem as a priority due to ageing society changes and introduced policy to improve health status among the older population. I found that in Eastern region still has inadequate research in this field in order to support the policy and improve ageing health.

In 2018, there was a survey conducted in Eastern of Thailand, which region is controlled by my work place. The survey was done in the topic of health behavior and healthy lifestyle among the elderly. I had attended to the research planning meeting and found it was interesting. The survey also covered the question of falls which was passionate me to study in this issue as it is a serious problem in ageing health and the incidence is high. The consequences of falls in elders are not only affect elders but also family members. Thus, I think the research in factors related to falls among the elderly will help to understand more in its cause and risk factors. The successful research, which result would benefit in fall prevention and reduce the incidence of falls among the elderly.

CHAPTER ONE : BACKGROUND OF THE COUNTRY

1.1 General information of Thailand and Eastern region

Thailand, is a country in the South-East Asia continent. The total area of the country is 513120 km² with a number of approximately 69.5 million total population in 2019(1). Thailand is divided into 6 geographic regions according to the Office of the Royal Society; North, South, East, North-East, West and Central, with a total of 77 provinces(2). The capital city, Bangkok, is located in the central region with approximately a population of 10 million (1). The official language is Thai and the major religion is Buddhist. Thai people have a common family-structure, which is described as that several generations live together in the same house. This means that older adults are more dependent when they usually stay with their children.

Thailand is considered as an upper-middle income country with a gross national 125,765 Thai baht per capita in 2019 (4210 Dollar US)(3). The economic growth in Thailand had been continued rising 5% annually, which brought many Thai people an improvement of their financial status. Poverty is still a concern in Thailand as same as inequity. The extensive increase of poverty was rising from 7.2% in 2015 to 9.8% in 2018(3), while the GINI index also increased from 36 to 36.4 in the same period(4). After the COVID-19 outbreak, affecting a decrease in tourism and trade, the financial status is expected to weaken in 2020.

The Eastern region of Thailand (figure 1) consist of seven provinces (figure 2); Chonburi, Rayong, Chachengsao, Chantaburi, Trad, Srakaew, Prachinburi, with a total area of 34,380 km² and a 4.7 million total population. This region has specific economic characteristics that are called the Eastern Economic Corridor (EEC) (see annex 1). The EEC, region in the eastern seaboard is covered by three provinces and is promoted to develop in industrial growth, including a decentralized economic growth(5).

Figure 1 The map of Eastern Thailand



Resource: Wikipedia website
Available from:

https://en.wikipedia.org/wiki/Eastern_Thailand

Figure 2 Provinces in Eastern Thailand



Resource: Thailand visitors website
Available from:


<http://thailandforvisitors.com/east/>

1.2 Demographic information

The literacy rate of the population age > 14 was 93% in the last update of 2015. The average population growth in Thailand is 0.4 % annually, and population age distribution among 0-14 years old is 16.8, while age 60 and above is 18.5 in 2019(1). In 2019 it was reported by Thai NSO, that 16.45% were in the 0-14 age group, 64.37% were in 15-59 age group and 16.73% were in the over 60 age group(6). The change of the Thai population structure is affected by a rising, ageing population which increased from 6.8% in 1994 to 14.9% in 2014, up to 16.7% in 2019(7).

The last update, in 2015, the fertility rate in Thailand was reported to be 1.5 live birth per women, which was a decrease from 2010. Life expectancy at birth increased to 78.9 in females and 71.6 in males in 2018, the crude birth rate was 9.6 per 1000 population, while the mortality rate was 7.1 per 1000 population(8).

Figure 3 Change in Crude Birth Rate, Mortality Rate and Natural Growth in Thailand 2018



Resource: The Bureau of Registration Administration, Department of Provincial Administration, Ministry of Interior

1.3 Health situation, health system and Public health concern

The major cause of death is non-communicable diseases (74%), which was reported in 2016. The highest mortality rate were cardiovascular diseases (23%) and other NCD (23%) followed by cancer (18%) (see annex 2). Other non-communicable diseases are considered as health problems, such as diabetes and hypertension, due to their prevalence have increased 3-4 times over 15 years(9).

The current health expenditure in Thailand in 2016 was 3.7% of the GDP(1). Universal health care is provided by a civic service welfare system for civil servants and their family, Social Security for private employees and universal coverage for all Thai nationals. The universal coverage was introduced since 2001 and it covers health care services for all the population, including low-income households, the user fee is 30 baht (0.96 US\$) per health care service but it usually covers basic treatment. The report from the World Bank has indicated that 99.5% of the total population of Thailand has health insurance coverage(10).


The health system in Thailand is controlled by the Thai Ministry of Public Health. Health care services are provided by public health sectors such as health posts in the community level, health centers, district hospitals, provincial hospitals. Private health sectors such as private hospitals and clinics are included in different system, the private hospital association.

CHAPTER TWO : PROBLEM STATEMENT AND METHODOLOGY

2.1 Problem statement

Nowadays, most countries around the world are experiencing the growth of the elderly in their countries, due to the development of the health system and the decrease in fertility rate. This demographic changing pattern shows not only the rising in aging of the population but also the falling of the proportion of young children (Figure 4), which causes a higher life expectancy globally(11). The double rise from 12% -22% of population aging globally is expected from 2015 until 2050, in which 80% of these are in low and middle-income countries(12). Thailand has senior citizens in 16.7 % of all population in 2019 (7) and is expected to become a super-aging society, 30 % of the population are at the age of 65 or above, by the next 20 years(13).

Figure 4 Young Children and Older People as a percentage of Global population,1950-5050




Source: United Nation. World Population Prospects:
The 2010 Revision
Available at: <http://esa.un.org/unpd/wpp>.

The degenerative process of the age changes leads to cognitive impairment, functional impairment and impairment of mobility, especially the last two issues can cause falls, this is one of the common geriatric syndromes, apart from urinary incontinence, pressure ulcers, delirium, and functional decline(14). According to the functional and mobility changes, they may have balance problems, muscle weakness, poor vision, or a long term health condition and cognitive changes such as dementia, which can lead to dizziness and loss of consciousness and fall (9.) Environmental is a part of the precipitate factors of falls, such as using stairs, slippery floors, or other risky structures of the house, that cause problems through their imbalance in movement. Especially when older adults fall, this may lead to many serious health issues, such as fracture, joint dislocation, head trauma, and fatality. Among the aging population, one of the most common injury causes of mortality is falls, which is a public health issue and is considered nowadays as a priority in the world health organization and the Thai Ministry of Health. Comparing with other age groups, falls in the elderly have a higher impact

on the mortality rate(16) (Figure 5) because of physical status, which degenerates by the aging change and including the presence of individual medical conditions.

Figure 5 The Mortality rate of falls Among the elderly and other age groups of the population from 2007-2016 in Thailand



Source: Division of Non-Communicable Disease, Thai Ministry of Health
 Available from: <http://www.thaincd.com/document/file/info/fall/1-elderly28122560.pdf>

According to statistical data in Thailand, the mortality rate caused by falling had continued rising from 8.5/100,000 population in 2011 up to 10.2/100,000 population in 2017 (16). Health region 6, consisting of 7 provinces in the Eastern area, including 1 province from central area Samutprakarn, experienced a high and increasing falls mortality rate, from 9.2 in 2011 to 11.3 in 2017, in which both numbers were above the average rate in the national report(16). According to the mortality rate of falls in Thailand, approximately 5 people died from falling per day, in which 2 of them were elders.

Falls in the elderly could cause a chance of hip fracture 1 in 10, and this situation will increase by age(17). One-fourth of the hip fracture cases in the elderly could lead to death. The consequences of falls cost money in terms of medical and personal care, even though less than half of the elderly attended health facilities afterward. The patients who experienced falls with hip fractures, or head traumas, will have less independence and mobility, and the pre-fracture function cannot recover. The negative fall affects, such as fear of falls, would remain with them, which will lead to a hypoactive lifestyle and functional decline(14). In case of incapacity to receive the operative treatment when needed, due to the high risk from the procedure, the result can be remaining bedridden, immobility, or inability to move. The outcome of these falls has an impact on living to the elderly themselves, to their caregivers, such as spouses and family members. Notably, the negative result from falls always needs physical, mental, and financial support from family members.

2.2 Justification

Numbers of research explored the factors of falls among the elderly globally; almost half of them are from western countries. Most of the studies have focused on the environmental factors of falls, such as living conditions and the characteristics of the house.

The movement disorders such as disorientated movement or muscle weakness and their causes, were also emphasized as the factors which had the attention in many researches. On the other hand, there are few studies who were exploring overall factors, including community or family support factors, that might affect falls in older adults, instead of focusing on only specific factors. There is no research in falls among the elderly, that has been conducted and studied about all the possible related factors in Eastern Thailand, where the mortality rate of falls has always been higher than the national average rate. This study area is mostly in the urban area and is filled with many industrial factories, which means older adults are left behind staying home, and some of them have responsibilities in taking care of their grandchildren instead of parents, who are fulltime workers.

One survey conducted in the East Thailand had the purpose to explore the related factors to health status and healthy lifestyle among the elderly, which consisted of general information of participants, such as weight, height, education, physical status, medical condition and life style. Besides, the questionnaire included the participants falls history, level of health literacy, housing, family support, and participating in community activities that is related to falls among the elderly. This study is planned to be published within next year, in it is explored its research question, which is the overall health status, healthy lifestyle and related factors between these issues among the elderly in the Eastern area, but did not explain the linkage specifically between falls and the factors that could link to it.

2.3 Research objective

2.3.1 General objective

To explore and analyze the factors associated with falls among the elderly in Eastern Thailand and to give recommendations regarding falls of elderly to Thai MoPH, health facilities, health workers and communities.

2.3.2 Specific objective

1. To explore and analyze individual (dependence and independence) causes of falls among the elderly in the Eastern Thailand
2. To explore and analyze factors in proximal cause-level that is related to falls among the elderly in the Eastern Thailand
3. To explore and analyze factors in distal cause that is related to falls among the elderly in the Eastern Thailand
4. To explore and analyze the best practices of falls' prevention in Thailand and other countries.
5. To give recommendations regarding falls of elderly to the Ministry of Health, health facilities, health workers and communities.

2.4 Methodology and analytical framework

This thesis is a mixed-method study that will be done by firstly, secondary data analysis and then literature review.

2.4.1 Secondary data analysis

2.4.1.a Study type

This method will be done by analyzing the secondary data, collected by the survey among the elderly in the Eastern Thailand in March 2018. The research “The related factors to health status and behaviour in elderly in the Eastern part of Thailand” belongs to Tippawan Jumpang, registered nurse and senior professional working at Health Center 6, Thailand. This was done with the purpose of exploring factors related to healthy lifestyle among the elderly in the health region 6.

The survey was conducted by collecting the data from face-to face-interview, which questions proved the content validity and realizability. Participants were selected by a multi-stage sampling method (see below). The weight, height and waist circumference were measured by the research team to calculate BMI and identify the normal value for analysing. Questions of incomes, physical status and medical conditions, as same as others questions, were collected from answers of participants. The research survey had been analyzed, the summary report was written and it is planned to be published in Thailand not later than 2021, with the original research heading.

The permission letter has been sent to the research owner for the authorization and the agreement (see annex 3), and it was replied before using the data. The waiver was submitted to the chair KIT REC with the reply of decision (see annex 4,5). The secondary data analysis is aimed to analyze and explore the specific objectives explained above. The analysis process will be done in each level of causes according to the analytical framework, to study the relation between various factors. The secondary data analysis will be done by using the SPSS program version 26.0 to find the descriptive data and to analyze statistically between two independent groups of data, falls and non-falls, with all factors in order to find the association. The collected data are categorical both nominal and ordinal and will be analyzed to find significant association ($p\text{-value} \leq 0.05$) by using the Chi-Square test, Fisher’s Exact Test will be used when the Chi-Square test cannot be done, and the odd ratio is used to find strength of the association.

2.4.1.b Study area

The study area is the health region 6, the eastern region including one province from the central region according to the regional health system in Thailand, which totally consists of 8 provinces.

2.4.1.c Study population

This study will be focused on the population in the communities that is older than 60 according to the WHO definition of an aging population. The analysis will be done from the survey “Related factors to health status and behaviour in elderly in health area 6 in Thailand”. The survey collected data from 450 elderly who live in health area 6. The number of participants were selected from total 512,857 elders in the Eastern part, by a three-stages cluster sampling method. Firstly, the simple sampling was done to select 3 from totally 8 provinces followed by a simple sampling method to select 3 districts from each province. And lastly, the simple sampling method was used to select 5 subdistricts from each district, in which only elders, who were physically active, were included in this research survey. If any subdistrict had less than 10 active elders, the participant from the neighbouring subdistrict would be included in the research survey.

2.4.2 Literature review

2.4.2.a Study type

The literature review in this thesis will be done according to a systematic approach, to explore all related factors from the published materials, both in the country and also in other countries. This method will be used as the causation conceptual framework, to analyze the factors related to falls among the elderly and to link and compare with the secondary data analysis. The literature review will be done by firstly reviewing the study which was done in Thailand and other South east Asian countries in terms of context, culture and social economic status similarity and then the research from other countries will also be included after that, in order to explore the specific objectives and comparing due to the background differences. Many literatures conducted in Western countries have the intervention in falls prevention, which will be reviewed as best practices in the specific objective.

2.4.2.b Search Strategy

The literature review method in this study used search engines: Google Scholar, VU online library, PubMed. The website of WHO, Thai Ministry of Health, NHS, Thai NSO were used to explore the definition, health information and health data. The searching process was done through Thai and the English language. Searching in the English language helped to find the international guideline and literature, which are done abroad or are published internally, but most of statistic data in Thai population and some Thai literature, needs to be explored in the Thai language. Keywords in MeSH term are falls in elderly, aged, gender, environmental hazardous, intervention, falls, community, senior housing, family support, elder abuse, policy, healthy lifestyle, medical condition, physical status. Articles were chosen according to the subjects followed by reviewing the abstracts, introduction, result, discussions and conclusion.

2.4.2.c Limitation of the methodology

The literature review had limited number of Thai studies within 20 years, while many literatures were done in Western countries, with a different social-economic status, context and culture. Literature review will be done by exploring international articles, to find factors in each level that have insufficient information and especially in best practice of intervention. The survey did not include fear of falls and violence among the elderly, which is one of the causes of falls among elderly, thus this factor will be studied by literature review method.

2.4.2.d Inclusion Criteria

All of the literatures, that are included in the review, were published less than 20 years in both Thai and the English language. Literature for the review was conducted in Thailand, South-East Asia countries and also other countries.

2.4.2.e Exclusion Criteria


The research was conducted and published more than 20 years ago will be excluded. The research in other languages except English and Thai will not be included in this study.

2.4.3 Analytical Framework

This research study uses the Causation of falls conceptual framework (Figure 6) which is adjusted from a causal-web in “Comparative quantification of health risks: conceptual framework and methodological issues”(18). The original framework from the survey was emphasized into three factors (individual, environmental and social security system) that are linked to healthy behaviours, which is impropriated with the research question of this study and covers not all specific objectives this research is aimed to explore. The Dahlgren-Whitehead model of determinant of health, seems to have various layers according to the determinant of health, but it was not used because the components of each factors were categorized differently from the purpose, and the problem tree of this study. The causation framework was chosen as reason of essential levels of causes of the outcome “Falls among the elderly” were well organized for understanding from the individual person to distal causes, community, the health system, policy and government. The structure of the causal-web is well illustrated, beside the different layers of the disease risks and causes, it showed the interaction within and between them.

This conceptual framework started from the outcome “Falls in elderly” and branched off its causes, which is divided into different levels of causes. The first level, the pre-proximal factor, is an individual factor, which means all causes of falls in this level are related to the elderly themselves and this can be dependence and independence. The individual dependent factors, refer to those factors that are relying on something or someone else such as age, gender, physical status and medical condition. On the other hand, independence refers to factors that are not controlled by other people or something such as lifestyle, level of health literacy. The proximal cause refers to environmental conditions which is the characteristic of the house and the area around the house. While family support, considered as proximal factors, will study the living condition of the elderly, whether they stay alone or with their families, include the support or abuse from family members. The distal cause refers to social and community factors, such as the support activities from a senior club, home visiting and violence issue. This conceptual framework considered the Government and policy as the distal cause, which might have an impact on the elderly subsistence allowance and the home visiting by health care workers and policy.

Figure 6 Causation of falls conceptual framework



CHAPTER THREE : STUDY FINDING

3.1 Finding introduction

In this chapter, the result of this study will be reported in 4 categories, including the background of the study finding, individual cause, proximal cause and distal cause. The result of descriptive data analysis and the prevalence of falls including causes and consequences among this study population will be explained in the background. This analysis excluded issues about dental care, dental check-up from the survey, as these are not related to falls and the objective of this study. This study aimed to find a relation between 5 major topics of healthy lifestyle, which will be explained in the independence individual part, and excludes other lifestyle topics from the survey: eating 3 meals, eating well cooked food, eating pickled fruits or vegetables, drinking soft drinks and eating betel nuts.

3.2 Description of the characteristics of respondents

The study reports from totally 450 participants this had 88 male participants and 362 female participants which is approximately the ratio of male to female 1:4. The age of participants ranged from 60 to 88 years old with the mean being 68.45 and SD= 5.852. The descriptive data showed percentage of male in the survey much less than in the regional statistics. The reason of this disproportion was explained by the researcher that the majority of male went out for work during the survey, which was working time. The same reason as in 60-69 age group had lower percentage and in the group age 80 and above had higher percentage comparing to the regional statistic data. Among all participants, 121 elders or 26.9 % were reported with having a history of falls within the 1-year period before the survey.


Table 1 Demographic data of study population among 450 respondents age >60 years old in a survey in Eastern Thailand in 2018

Characteristic		N	%	N, % of population age >60 in Eastern Thailand in 2019 (7) (N total= 719044)
Gender	Male	88	19.6	315119, 43.8%
	Female	362	80.4	403925, 56.2%
Age-group	60-69 yrs	285	63.3	402609, 56%
	70-79 yrs	142	31.6	203451, 28.3%
	>80 yrs	23	5.1	112984, 15.7%
Falls	Have falls history	121	26.9	
	No falls history	329	73.1	

3.2.1 Falls among respondents

Causes of falls in the research survey were different (Figure 7), the most frequent cause of falls was stumbling, followed by slipping and other causes, which were 41.6% and 26.4% and 11.7%. Other causes indicated in this study were gaps or steps on the floor, during biking, miss stepping, falling from the stair, during physical activities, and muscle sparking. Apart from the majority of causes, dizziness leads to falls 6.7% and muscle weakness resulted in falls 5%.

Figure 7 Causes of falls pie chart, among 121 respondents age >60 years old who had fallen in the past year in a survey in Eastern Thailand 2018



Among 121 respondents who experienced falls, 71 elders fell inside the house and 53 elders fell outdoors, while 3 among them had fallen both at home and outside the house. The most frequent falls cases inside the house was on the balcony, which is 27.1% followed by other places 22.8% such as at the door, the corridor and the area around the house, while falls occurred at the stairs was reported as the smallest proportion in this study.

The survey had focused temples and hospitals as places of fall outside the house, because these places are considered older adults usually go to in the Thai context. There were only 3 people that had fallen at temples and no one had fallen in hospitals. Other places outside the house, such as the field and the rubber plantation, were also indicated in this survey.

3.2.2 Consequences of falls among respondents

The consequences of falls were a majority injured (n=88) and only 33 among them were non-injured. The most severe result was a bone fracture, which occurred in 6 elders and 4 elders had head laceration wounds. The minor symptoms are contusion (n=49), pain (n=21) and abrasion wounds(n=18). Besides, in the survey 10 elders were recorded with other symptoms such as hematoma, hip joint displacement, tendon injury and teeth injury. Although three fourth of falls cases were injured but only one-third (n=42) visited the doctor and 6 elders among those who fell, had to be admitted to hospital. While 34 fallers didn't have any treatment and 45 fallers were self-cured.

3.3 Individual cause

3.3.1 Dependence factors

3.3.1.a Age

The participants in this study were 60 years and older, I have grouped them into a 3 age-group with a 10 years interval, according to the guideline of Foundation of Thai Gerontology Research and the Development Institute(13). More than half of the participants were in the 60-69 years old age-group and the incidence of falls in this groups was the lowest

(n=68, 23.9%) comparing to those who did not fall, while eight from the 23 respondents age over 80, had the highest percentage of falls (34.8%). This study illustrated that the older age-group fell more, although no significant association was reported (see table 3).

There is a cross-sectional study (n=26689) from the national survey of Sophonratanapokin et al that reported the effect of older age to an increasing in chance of falls(19). Likewise, the study of Worapanwisit et al (n=406) in a Southern province had described the correlation of falls with older age(20). On the other hand, the study in a senior house in Bangkok of Maneeprom et al did not find that older age was a risk factor of falls among the elderly, while this research (n=64) included physically active participants without dementia and hearing impairment, which may be selection bias(21).

3.3.1.b Gender

The result of the study found among male, 19 of them (21.6%) fell, while female fell more (n=102, 28.2%). Females seem to have more effect in falls in the survey, the result of analysis showed no significant association between gender and falls cases (see table 3).

Several Thai articles explained differently, the research of Thai national survey explained that females were likely to fall more than males with a significant association (19). In the same ways, Chaiwanichsiri et al observed the adjusted odds of falls among women, were 2.3 times greater than men in the study (n=213) in urban Bangkok. In their study used potential factors to analyse adjusted Odds Ratio, which were foot pain, pes plantar, knee OA(22).

3.3.1.c Education, Marital status, Religion, occupation and financial status

Table 2 Descriptive data of education, marital status, religion, occupation and financial status among total 450 participants age >60 in the survey in Eastern of Thailand in 2018

Characteristic	Number	Percentage	
Marital status	Single	25	5.6
	Marriage	227	50.4
	Widow	148	32.9
	Divorce	50	11.1
Religion	Buddhist	425	94.4
	Christian	1	0.2
	Muslim	24	5.3
Education	No education	15	3.3
	Primary school	333	74.0
	Secondary school	38	8.4
	High school	35	7.8
	Above high school	29	6.4
Occupation	Not working	223	49.6
	Farmer	55	12.2
	Handyman	95	21.1
	Own business	1	0.2
	Officer	61	12.6

	Others	15	3.3
Source of incomes	From work	219	48.6
	From Family	254	56.4
	From Elderly allowance	406	90.2
	From pension	23	5.1
Sufficiency of income	Sufficient	304	67.6
	Not sufficient	146	32.4

This study aimed to identify the relation between education level and falls, firstly to analyze those who studied and who did not, which results showed no effect of educated and non-educated to falls. Analyzing further between falls and different level of education, the result illustrated participants who had the highest level of education fell most (n=9, 31%), participants who did not fell fewer (n=4, 26.7%) and there was no report of significant association between them (see table 3).

In the same way, the literature of Maneeprom et al, in which a study in a senior house in Bangkok(n=64) found no association between falls and levels of education(21). There were a number of articles in Thailand that showed a low level of education effecting on the increasing falls incidence among the elderly with the report of a significant association. The study (n=406) of Worapanwisit et al in a Southern province illustrated elders, who finished primary school, fell most, while older adults who had a bachelor degree fell less (20). In the same way, the large sample size study, in the national survey, observe that the highest proportion of their respondents who fell did not study, while the smallest proportion was among who graduated from colleges or had higher degrees(19).

The result from the survey showed different marital status, religion and occupation were not related to falls, where the statistically analysis reported no significant association. To explore the relation of career and falls, this study was aimed to find association between working and non-working groups. Although participants who worked fell slightly more than who did not work, no significant association was reported in the result of analysis. I did not find other literature in Thailand that explored these factors to falls.

In this survey the observed participant, who had insufficient income, fell (n=47, 32.2%) than who had sufficient income (n=74, 24.3%), this study reported no significant association between them (see table 3). The result from the survey defined sufficient income by participants' answers, which may be an information bias. While the study from national survey (n=26689), found insufficient income affecting falls. This study used the report of incomes but did not explain the cut-off point in their article. After adjusting potential factors, his study showed the odds of falls among elders, who had insufficient income, were 1.39 times greater than who had sufficient (95% CI= 1.28-1.51). Potential factors that were used in this study were living environment, gender, health status, health behavioural, education and location of home(19).

Table 3 Bivariate analysis of dependent individual factors with falls in a survey in Eastern Thailand 2018

Characteristic	Falls within 1 year: N (% ^a)		p-value	OR	95% CI
	Yes	No			

Age (years old)			0.155		
- 60-69	68 (23.9)	217 (76.1)			
- 70-79	45 (31.7)	97 (68.3)			
- 80-89	8 (34.8)	15 (65.2)			
Gender			0.211		
- Male	19 (21.6)	69 (78.4)		0.7	0.40-1.23
- Female	102 (28.2)	260 (71.8)		1	
Education			0.780		
- No education	4 (26.7)	11 (73.3)		0.99	0.31-3.17
- Literate	117 (26.9)	318 (73.1)		1	
• Primary school	92 (27.6)	241 (72.4)			
• Secondary school	7 (18.4)	31 (81.6)			
• High school	9 (25.7)	26 (74.3)			
• Above high school	9 (31)	20 (69)			
Marital status			0.108		
- Single	4 (16)	21 (84)			
- Married (stay together)	59 (26)	168 (74)			
- Widow	38 (25.7)	110 (74.3)			
- Divorce	20 (40)	30 (60)			
Religion			1.000 ^c		
- Buddhist	115 (27.1)	310 (72.9)			
- Christian	0 (0)	1 (100)			
- Muslim	6 (25)	18 (75)			
Occupation			0.749 ^c		
- Not working	63 (28.3)	160 (71.7)		1.15	0.76-1.74
- Working	58 (25.6)	169 (74.4)		1	
• Handyman	12 (21.8)	43 (78.2)			
• Farmer	24 (25.3)	71 (74.7)			
• Officer	0 (0)	1 (100)			
• Business owner	16 (26.2)	45 (73.8)			
• Others	6 (40)	9 (60)			
Sufficient incomes:			0.079		
- Yes	74 (24.3)	230 (75.7)		0.678	0.44-1.05
- No	47 (32.2)	99 (67.8)		1	

^a % of sub-group in each characteristics (in the row)

^b p-value<0.05: significant association

^c Result was analyzed by using Fisher's Exact Test

3.3.1.d Physical status (BMI, waist circumference, visual impairment, hearing impairment, balancing impairment, taking sleeping pills)

Table 4 Descriptive data of Physical and health status among total 450 participants age >60 in the survey in Eastern of Thailand in 2018

Characteristic	n	%	
BMI ¹	Underweight	12	2.7
	Normal	137	30.4
	Overweight	80	17.8
	Obesity	178	39.6
	Severe Obesity	43	9.6
Abnormal waist circumference	339	75.3	
Visual impairment	183	40.7	
Hearing impairment	67	14.9	
Balance impairment	105	23.3	
Taking sleeping pills	207	46.0	
History of hospitalization within 1 year	62	13.8	
Health checking up within 1 year	382	84.9	

This study had categorized the level of BMI into 5 groups according to the Thai MoPH guideline, which consists of underweight (BMI<18.5), normal (BMI=18.5-22.9), overweight (BMI=23-24.9), obesity (BMI=25-29.9) and severe obesity (BMI>30). The result illustrated elders with a normal BMI fell 19% (n=26), while the overweight (n=19, 23.8%) and obese group (n=56, 31.5%) fell more and the largest proportion (n=17, 39.5%) was in the severe obese group. This study well explains the increasing chance of falls in the higher BMI participants and there was a significant association between BMI and falls (see table 6). In addition, older adults who had a low BMI are more likely to fall than normal BMI elders. This study had explored the relation of waist circumference, which is associated to BMI, to falls. The abnormal waist circumference in a Thai male is more than 90 centimetres and more than 80 centimetres for the female, according to the guideline of Thai MoPH. The result of this study found participants who had abnormal value, fell more (n=108, 31.9%) than who had a normal value (n=13, 11.7%) with the evidence strong association (OR=3.52) between having abnormal waist circumference and falls (see table 6).

The study of the physical status aimed to find the effect of visual, hearing, balance impairment and taking sleeping pills, to falls. The result reported respondents with visual impairment fell slightly more than those who had normal vision, however, there was no evidence of significant association. This study found that hearing impairment, balance impairment, and taking sleeping pills had the effect to falls with report of significant association (see table 6). Among these factors, balance impairment was found strongly associated to falls and it can be explained that the odds of falls among participants who had poor balance, were greater almost three time than who had normal balance (see table 6). This survey also explored the history of admission in the hospital and health check-up within 1 year were no significant association in these issues with falls were observed.

Several Thai literatures had studied the effect of physical status among the elderly to falls. The study (n=406) in a Southern province reported the cases of falls, were higher in overweight and obese participants, with a statistically significant association. The same study found elders who had visual problems, fell more with a significant association but their study

¹ BMI = kg/m²

focused on participants of a narrower age group (60-69 years old)(20). As same as the study (n=213) in urban Bangkok, it reported that poor vision strongly related to the increased risk of falls. Especially in females, this study found the odds of falls among female elders, who had visual impairment, were almost 5 times greater than females who had normal vision (OR=4.7)(22).

The cross-sectional study between falls and physical status (n=90) of the elderly, in a rural community, aimed to find the physical ability, by using four physical status measurements, which results illustrated poor physical ability is the risk factor to falls among the elderly. This study only included participants age 75 or above, BMI 18.5-29.9, and without neuromuscular disease to avoid the walking ability effect but the these may be selection bias (23).

3.3.1.e Medical conditions (diabetes mellitus, hypertension, ischemic heart disease, dyslipidemia, depression, cancer, knee osteoarthritis)

Table 5 Descriptive data of medical conditions among total 450 participants age >60 in the survey in Eastern of Thailand in 2018

Characteristic		n	%
Medical condition	Diabetes mellitus	115	25.6
	Hypertension	240	53.3
	Dyslipidemia	179	39.8
	Ischemic heart disease	35	7.8
	Depression	3	0.7
	Cancer	3	0.7
	Knee osteoarthritis	75	16.7

This study aimed to find the relation between falls and medical conditions, which are diabetes mellitus (DM), hypertension (HT), ischemic heart disease (IHD), dyslipidemia (DLP), depression, cancer and knee osteoarthritis (knee OA). The data were collected from answers of participant without the evidences of medical record, which may be a measurement bias. Each of the diseases were divided into participants who suffered from disease and those who did not sufferer, including who did not know.

The result showed that 33 of DM participants (28.7%) experienced falls and non-DM participants had a slightly lower percentage of falls (n=88, 26.3%). In the same way, participants with HT fell (n=73, 30.4%) more often than who did not suffer from HT. Among 35 participants with IHD, 12 of them fell (34.3%) which was higher than those who did not suffer from IHD (n=109, 26.3%). Participants who had DLP had fallen similarly to those who did not suffer from this disease.

The medical conditions, which are not lifestyle related, were rare among respondents and significant associations were not identified. Only three participants were diagnosed with depression and one of them had a fall history, while participants without depression fell less (26.8%). Among three participants with cancer, two of them had a fall history, while the group with non-cancer participants, experienced falls much less (26.6%). The result showed that participants with knee osteoarthritis (knee OA) fell (36%) more often than participants without knee OA (25.1%). Although this study observed the higher incidence of falls among DM, HT, IHD, DLP, depression, cancer elders, but there was no report of significant association

between them. Knee OA was found not associated with fall while the result was just above the border line of significant association (see table 6).

Table 6 Bivariate analysis of dependent individual factors with falls in a survey in Eastern Thailand 2018

Characteristic	Falls (% ^a)	None-falls (% ^a)	OR	95% CI	p-value
BMI					0.038 ^b Cramer's V=0.15
- Underweight	3 (25)	9 (75)			
- Normal	26 (19)	111 (81)			
- Overweight	19 (23.8)	61 (76.3)			
- Obesity	56 (31.5)	122 (68.5)			
- Severe Obesity	17 (39.5)	26 (60.5)			
Excess waist circumference			3.52	1.89-6.57	<0.001 ^b
- Yes	108 (31.9)	231 (68.1)			
- No	13 (11.7)	98 (88.3)			
Visual impairment			1.31	0.86-1.99	0.210
- Yes	55 (30.1)	128 (69.9)			
- No	66(24.7)	201 (75.3)			
Hearing impairment			1.78	1.03- 3.07	0.037 ^b
- Yes	25 (37.3)	42 (62.7)			
- No	96 (25.1)	287 (74.9)			
Balance impairment			2.97	1.87-4.71	<0.00 ^b
- Yes	47 (44.8)	58 (55.2)			
- No	74 (21.4)	271 (78.6)			
Taking sleeping pills			1.60	1.05-2.43	0.027 ^b
- Yes	66 (31.9)	141 (68.1)			
- No	55 (22.6)	188 (77.4)			
Diabetes mellites			1.13	0.71-1.81	0.613
- Yes	33 (28.7)	82 (71.3)			
- No	88 (26.3)	247 (73.7)			
Hypertension			1.48	0.97-2.25	0.071
- Yes	73 (30.4)	167 (69.6)			
- No	48 (22.9)	162 (77.1)			
Ischemic heart disease			1.47	0.71-3.04	0.304
- Yes	12 (34.3)	23 (65.7)			
- No	109 (26.3)	306 (73.7)			
Dyslipidemia			1.04	0.68-1.59	0.850
- Yes	49 (27.4)	130 (72.6)			
- No	72 (26.6)	199 (73.4)			
Depression			1.36	0.12-15.16	1.000 ^c
- Yes	1 (33.3)	2 (66.7)			
- No	120 (26.8)	327 (73.2)			
Cancer			5.51	0.50-61.35	0.177 ^c
- Yes	2 (66.7)	1 (33.3)			
- No	119 (26.6)	328 (73.4)			

OA knee					
- Yes	27 (36)	48 (64)	1.68	0.99-2.85	0.051
- No	94 (25.1)	281 (74.9)			
History of admission in the hospital within 1 year					0.051
- Yes	23 (37.1)	39 (62.9)	1.75	0.99-3.07	
- No	98 (25.3)	290 (74.7)			
Health checking up within 1 year					0.498
- Yes	105 (27.5)	277 (72.5)	0.81	0.44-1.48	
- No	16 (23.5)	52 (76.5)			

^a % of sub-group in each characteristics (in the row)

^b p-value<0.05: significant association

^c Result was analyzed by using Fisher's Exact Test

While the analysis from the survey shows non-association result of knee OA, the study of Chaiwanichsiri (n=213) et al in urban Bangkok, found a strong association between knee OA and falls. After adjusted potential factors, this study reported the odds of falls among elders with knee OA, were 3.2 times greater than those who did not suffer knee OA. The potential factors were gender, foot pain, and pes planus. This study was done in the hospital, which may be a selection bias. The same study explored other lower extremities diseases, which observed foot pain (OR=2.5) and plantar fasciitis (OR=6.8) which were strongly associated to falls(22), although the survey did not focus on these conditions. A case control study (n=333) of Kuhirunyaratn et al, conducted in the province in North-East Thailand, explains the effect of depression to falls. They reported a significant association after adjusted the potential confounding between falls and depression (AOR=1.76, 95%CI=1.03- 2.99), this study did not indicate which confounding they used for analysing (24).

Besides, a descriptive study of Muangpaisan et al (n=80) explained falls occurred most among patients with a neurological disorder (Parkinson's disease and dementia) 40%, followed by musculoskeletal disorder (28.8%), visual impairment (18.8%) and postural hypotension (12.5%). This study only included hip fracture patients, which can be selection bias, but the results can explain more the medical condition cause, that leads to severe falls(25).

3.3.2 Independence factors

3.3.2.a Healthy Lifestyle (eating vegetable, drink water, exercise, not smoking, not drinking alcohol)

Table 7 Descriptive data of healthy life style among total 450 participants age >60 in the survey in Eastern of Thailand in 2018

Characteristic	N	%	
Eating fruits and vegetable regularly	177	39.3	
Drink water 8 cups/day regularly	304	67.6	
Exercise regularly			
	15-30minutes	289	64.2
	Balancing exercise	33	7.3

	Waving arms and legs	247	54.9
Not smoking		430	95.7
Not drinking alcohol		420	93.3

The Thai MoPH described the definition of a healthy lifestyle in the elderly, which contains 5 essential issues: 1. Doing exercise for 30 minutes not less than 5 days a week, 2. Eating fresh vegetables and fruits regularly, 3. Drink enough water at least 8 cups daily, 4. Not smoking and 5. Not consuming alcohol drinks. These are also included in the Key Performance Indicators among the aging population of Thai MoPH, to achieve the health national target(26). The guideline identified the frequency of each topic 5-7 days were concerned as healthy lifestyle, this study categorized participants who did the regular activity 5-7 days per week and those who did less than that will be concerned as unregular.

In general, the results showed slightly lower percentages of falls in the groups with healthy behaviours, compared to those without, however none of those percentages differed significantly to those without healthy behaviours. The result showed, among participants who ate vegetable and fruits regularly, fell less than who did not, as same as participants who regularly drank 8 cups of water per day, fell less than who did not. Most of the participants in this study did not drink alcohol and among them (n=115, 27.4%) had experienced falls, which is higher than those who drink alcohol (n=6, 20%). Participants who did not smoke fell (n=113, 26.3%) much lower than who smoke (n=8, 40%). No significant association between smoking/drinking alcohol and falls was observed in the survey (see table 8).

The analysis in physical activity, found elders who did exercise regularly, fell less (n=75,26%), although no significant association was reported. This study also emphasized two different types of exercise in which the first is waving arms and legs, one of the exercises recommended by the Thai Health Promotion Foundation(27) for the elderly(28). The second type is a balancing exercise such as yoga and Tai Chi, which is recommended for muscle strengthening for elders(28). This study illustrated that the two types of exercise did not affect the falls incidence among the elderly.

Several research conducted in Thailand, aimed to find the effect of healthy behaviours to falls, among the elderly. The large sample size study from the national survey, reported a similar result of no significant association between the regular intake of vegetables, alcohol consumption and falls(19). The same study reported respondents, who did not do regular exercise, fell more than who did with a significant association(19). Similarly results of a medium sample size case control study (n=333), showed that doing sufficient exercise, was the protective factor to falls and it was significantly associated(24). I did not find other articles looking into other healthy lifestyles and falls in Thailand.

3.3.2.b Level of health literacy

The level of health literacy (HL) refers to the ability of an individual to access, understand, ask, share, exchange the health information with others and behavioural/lifestyle change to obtain good health. This study had analyzed 10 survey questions and was categorized into two levels, which are low (less than 50%) and high (more than 50%). The result of this study showed that the group with a high level of HL fell more. A similar result, when this study grouped HL into 3 levels, which were <50%= Inadequate, 50-80%= Good, >80%= Excellent, there was no significant association reported. Although the HL

level was not related to falls in this survey, I did not find other researched explored effects of HL to falls (see table 8).

Table 8 Bivariate analysis of independent individual factors with falls in a survey in Eastern Thailand in 2018

Characteristic	Falls (% ^a)	None-falls (% ^a)	OR	95% CI	p-value
Eating fruits and vegetable regularly					0.260
- Yes	33 (23.4)	108 (76.6)	0.767	0.484-1.217	
- No	88 (28.5)	221 (71.5)			
Drink enough water regularly					0.866
- Yes	81 (26.6)	223 (73.4)	0.963	0.618-1.500	
- No	40 (27.4)	106 (72.6)			
Exercise 15-30 minutes regularly					0.548
- Yes	75 (26)	214 (74)	0.876	0.569-1.349	
- No	46 (28.6)	115 (71.4)			
Balancing exercise					0.708
- Yes	22 (25.3)	65 (74.7)	0.903	0.528-1.542	
- No	99 (27.3)	264 (72.7)			
Waving arms and legs exercise					0.902
- Yes	101 (27)	273 (73)	1.036	0.592-1.812	
- No	20 (26.3)	56 (73.7)			
Not smoking					0.176
- Yes	113 (26.3)	317 (73.7)	0.535	0.214-1.342	
- No	8 (40)	12 (60)			
Not drinking alcohol					0.378
- Yes	115 (27.4)	305 (72.6)	1.508	0.601-3.784	
- No	6 (20)	24 (80)			
Level of Health Literacy					0.133
- High	79 (29.5)	189 (70.5)	0.718	0.465-1.107	
- Low	42 (23.1)	140 (76.9)			

^a % of sub-group in each characteristics (in the row)

^b p-value<0.05: significant association

^c Result was analyzed by using Fisher's Exact Test

3.3.2.c Fear of falls

The survey did not explore the issue of fear of falls, although it could relate to falls in both risk factor and consequences. Fear of falls can cause restriction in physical activities, and finally can end up with a fall from the effect of gait change(29). The cross-sectional study (n=386) was conducted to explore the fear of falls and its factors in elderly in Southern Thailand. The result showed half of both fallers and non-fallers were afraid to fall. The study

also illustrated perceived general health, visual impairment, mobility impairment, balance impairment, stroke, hypertension, antihypertensive medication, number of medication, and daily activity had an effect on this feeling(30).

3.4 Proximal cause

3.4.1 Environmental factor

Table 9 Descriptive data of living condition and area around houses of total 450 participants age >60 in the survey in Eastern of Thailand in 2018

Characteristic		n	%
House	One-story	196	43.6
	One-story with raised floor	86	19.1
	Two-stories	132	29.3
	Row wooden	3	0.7
	Town-house	22	4.9
	Others	11	2.4
Using stair	No	266	59.1
	Yes		
	> 5 times/day	75	16.7
	3-5 times/day	52	11.6
	1-2 times/day	57	12.7
Having handrails		210	46.7
Toilet	Squat	98	21.8
	Seat	308	68.4
	Both	44	9.8
Lightening	Sufficient	440	97.8
Area outside the house	Wet pathway with stagnant water	12	2.7
	Slippery ground	12	2.7
	The pothole in the ground	8	1.8
	Rough ground with rocks	33	7.3
	With obstacles	9	2
	Clear area	89	19.8
	With discarded items around	65	14.4
	With broken item around	24	5.3
	Inclined pathway	18	4

The study in the characteristic of the houses, showed the highest percentage of falls was observed in participants who lived in wooden row houses, followed by two-story houses. Elders who live in one-story housed fell more than elders who live in one-story house with

raised floors, although this type of house might have more environmental risk. No significant association between characteristic of the houses and falls was found in this study.

To compare the using of stairs and none-using stairs participants, the result showed that none-using stairs elders fell slightly less (n=71, 26.7%) than the using stairs participants (n=50, 27.2%) but no significant association between them was found.

This study observed having handrails to support their movement in houses not related to falls, due to similar percentages of falls cases in the group with handrails and without. Types of toilets were not associated to falls in the survey. The result showed the highest percentage of falls in those who use seat toilets (n=86, 27.9%), while using squat toilets, might cause more imbalance movement for the elderly, in contrast it had a lower percentage of falls (n=25,25.5%). Type of toilet has a relation to the financial status when a seat toilet is more likely to be found in a new house or in the urban area.

The study also emphasized different characteristics of the toilet floor; coarse, rough and slippery surfaces. Among elders, who had smooth surface toilets fell most (n=27, 37.5%) while who had rough and coarse surfaces of toilet fell less (n=90,25.1%; n=4, 20%). This study explained the higher cases of falls occurring in the slippery toilet, no statistically significant association was reported in this study. The association between the lighting and falls in the survey was not found, although elders who had dim light in their houses fell much more (n=4, 40%) than those who had sufficient light (n=117, 26.6%) (see table 10).

Table 10 Bivariate analysis of indoor environmental factors with falls in a survey in Eastern Thailand 2018

Characteristic	Falls (% ^a)	None-falls (% ^a)	OR	95% CI	p-value
Characteristic of the house					0.553 ^c
- One-story with raised floor	21 (24.4)	65 (75.6)			
- Two- stories house	40 (30.3)	92 (69.7)			
- Row wooden house	1 (33.3)	2 (66.7)			
- Townhouse	5 (22.7)	17 (77.3)			
- One-story house	49 (25)	147 (75)			
- Others	5 (45.5)	6 (54.5)			
Using stair					0.910
- Yes	50 (27.2)	134(72.8)	1		
- No	71 (26.7)	195 (73.3)	0.976	0.64- 1.49	
Having Handrails in the House					0.921
- Yes	56 (26.7)	154 (73.3)	1		

- No	65 (27.1)	175 (72.9)	1.02	0.67-1.16	
Type of Toilets					0.723
- Squat toilet	25 (25.5)	73 (74.5)			
- Seat toilet	86 (27.9)	222 (72.1)			
- Both	10 (22.7)	34 (77.3)			
Surface of toilet floor					0.076
- Coarse surface	4 (20)	16 (80)			
- Rough surface	90 (25.1)	268 (74.9)			
- Slippery surface	27 (37.5)	45 (62.5)			
Lightening					0.469 ^c
- Sufficient light	117 (26.6)	323 (73.4)	0.543	0.15-1.96	
- Insufficient light	4 (40)	6 (60)			

^a % of sub-group in each characteristics (in the row)

^b p-value<0.05: significant association

^c Result was analyzed by using Fisher's Exact Test

The hazardous outdoor environment was found a risk factor to falls in the survey, which results showed that areas with rocks, obstacles, discarded item, non-clear area and inclined pathways, affect falls among the elderly. The result showed that only a few participants had wet pathways (n=3), slippery ground (n=3) and potholes (n=2) around their house, which result recorded a similar percentage of falls among who had and who had not this hazardous situation with no significant association was reported. Among participants who had rocks in their pathway, 14 of them falls (42.4%), while participants who had a pathway without rocks fell less (n=107,25.7%). The result showed significant association between falls and a rough pathway with rocks which can be explained that the odds of falling among participants, who had rocks in their pathways, were 2.14 times higher than who did not (see table 11).

The majority of participants had no obstacles around their houses (n=117) and approximately one-fourth of them fell. While four from nine participants with barricades around their house fell, although no significant association was reported. This study is also aimed to explore the connection between the clear area around houses of participants and falls, which results showed 33 elders among who had a non-clear area fell (37.1%), which proportion was higher than who had a clear area (n=88,24.4%). The result showed a significant association and explained the odds of falls among participant who had an unclear area around their houses were 1.83 times higher than the other group (see table 11). The participants who had discarded items around their houses had fallen 36.9% (n=24), while another group fell less (n=97, 25.2%). The result by Chi square test showed a border line significant association but it was not relevant when Odds Ratio was calculated (see table 11). The result of this variable was uncertain, it may be affected by the sample size.

One-fourth of participants who did not have broken things around their houses, fell (n=116), while the other group fell more but no significant association was reported. The

study in a sloped pathway in front of the elders' houses found more participants who had this fell more (n=9, 50%) than who did not (n=112, 25.9%). The results showed the odds of falls among participants who had an inclined pathway were almost 3 times greater than those who did not have (see table 11).

Table 11 Bivariate analysis of outdoor environmental factors with falls in a survey in Eastern Thailand 2018

Characteristic	Falls (% ^a)	None-falls (% ^a)	OR	95% CI	p-value
Wet pathway with stagnant water					1.000 ^c
- Yes	3 (25)	9 (75)	0.904	0.241-3.396	
- No	118 (26.9)	320 (73.1)	1		
Slippery ground					1.000 ^c
- Yes	3 (25)	9 (75)	0.904	0.241-3.396	
- No	118 (26.9)	320 (73.1)	1		
The pothole on the ground					1.000 ^c
- Yes	2 (25)	6 (75)	0.905	0.180-4.545	
- No	119 (26.9)	323 (73.1)	1		
Rough ground with rocks					0.037 ^b
- Yes	14 (42.4)	19 (57.6)	2.135	1.034-4.405	
- No	107 (25.7)	310 (74.3)	1		
With obstacles					0.259 ^c
- Yes	4 (44.4)	5 (55.6)	2.215	0.585-8.390	
- No	117 (26.5)	324 (73.5)	1		
Clear area					0.015 ^b
- Yes	88 (24.4)	273 (75.6)	1,828	1.117-2.992	
- No	33 (37.1)	56 (62.9)	1		
With discarded items around					0.049 ^b
- Yes	24 (36.9)	41 (63.1)	1.738	0.999-3.024	
- No	97 (25.2)	288 (74.8)	1		
With broken item around					0.464
- Yes	8 (33.3)	16 (66.7)	1.385	0.577-3.324	
- No	113 (26.5)	313 (73.5)	1		
Incline pathway					0.031 ^{b,c}
- Yes	9 (50)	9 (50)	2.857	1.106-7.378	
- No	112 (25.9)	320 (74.1)	1		

^a % of sub-group in each characteristics (in the row)

^b p-value<0.05: significant association

^c Result was analyzed by using Fisher's Exact Test

The data analysis observed that the indoor environment did not affect falls, while outdoor environmental hazards were falls risk factors. Literature review found a study

(n=406) of Worapanwisit et al in a Southern province that reported that participants fall outdoor more, and the most common place was the area around their houses, with the evidence of significant association. The kitchen was reported as the most frequent place of falling indoors, followed by the bedroom, and living room with a statistically significant association. This study found the home hazardous environment such as contact with other persons or objects, slippery floor and slope way, affect the incidence of falls among the elderly(20).

A similar result from a large sample size research in the national level of Sophonratanapokin et al observed an increasing incidence of falls related to a slippery first floor and slippery bathroom. (19). The case control study (n=333), also reported a slippery floor is significantly associated to falls, and toilets without handrail were observed as a risk of falls among the elderly (24). The survey and Thai literatures did not find the effect of insufficient light. Poor lighting was reported significant associated to falls in several international studies. Dim light inside the house, especially on the stairs, lead to poor vision during the movement and can lead to falls(31)(32).

Descriptive analysis (n=80) of Muangpaisan et al was done among fall patients who were admitted with hip fractures, documented 78.6% happened indoors, and the most common places were bedrooms and bathrooms. Slippery floor and multi-stories floors, were reported as the most common cause. Only 12.5% of elderly in this study fell outside the house and it most frequent occurred in front of the house, but there was an unspecific character of this area(25). Although this study had a selection bias due to that only patients with hip fractures were included, the result illustrated more likely the factors that lead to severe outcome of falls.

3.4.2 Family factors

Table 12 Descriptive data of living condition and family support of total 450 participants age >60 in the survey in Eastern of Thailand in 2018

Characteristic		N	%
Living condition	Alone	43	9.6
	With spouses	80	17.8
	With children	322	71.6
	Other (parents, relative)	5	1.1
Major source of income	Elderly themselves	223	49.6
	Spouses	27	43.3
	Children	195	6
	Relative	5	1.1
Main care giver	Elderly themselves	82	18.2
	Spouses	76	16.9
	Children	274	60.9
	Relative	17	3.8
	Neighbour	1	0.2

3.4.2.a Family support

Living condition and family support were explored in secondary data analysis in order to find their effect to falls. The result illustrated 83 elders who live with their children fell

(25.8%), which incidence was lower than participants who lived with spouses (n=23, 28.7%). Elders who lived alone fell the most (n=15, 34.9%), which showed a higher percentage is among this group. No significant association between living with family and falls was reported. This study aimed to find a relation between source of majority income of the elderly, which result showed these variables were not associated to falls. In the same way, the result observed that the main care giver had no impact on falls (see table 13).

Table 13 Bivariate analysis of proximal factors with falls in a survey in Eastern Thailand 2018

Variables	Falls (% ^a)	None-falls (% ^a)	p-value
Living condition			0.350 ^c
- Alone	15 (34.9)	28 (65.1)	
- With spouses	23 (28.7)	57 (71.3)	
- With children	83 (25.8)	239 (74.2)	
- Other (parents, relative)	0 (0)	5 (100)	
Major source of income			0.971 ^c
- Elderly themselves	59 (26.5)	164 (73.5)	
- Spouses	8 (29.6)	19 (70.4)	
- Children	53 (27.2)	142 (72.8)	
- Relative	1 (20)	4 (80)	
Main care giver			0.889 ^c
- Elderly themselves	24 (29.3)	58 (70.7)	
- Spouses	20 (26.3)	56 (73.7)	
- Children	74 (27)	200 (73)	
- Relative	3 (17.6)	14 (82.4)	
- Neighbor	0 (0)	1 (100)	

^a % of sub-group in each characteristics (in the row)

^b p-value<0.05: significant association

^c Result was analyzed by using Fisher's Exact Test

Literature review found the study of Sophonratanapokin et al with a large sample size that explained the elderly, who lived with spouses, fell less than who lived alone with a statistically significant association. This can be explained that living with spouses is the protective factor to falls among the elderly (OR= 0.68, 95%CI=0.59-0.78)(19).

3.4.2.b Violence from family members

Although the research survey did not explore the violence issue among the elderly in particular area, there were a few studies of elder abuse that reported a high prevalence (78.4%) one in the Eastern province and 46.4% in female elderly in Northern province(33). Another study explored the relationship between violence among female elders and family factors. The study (n=304) of Chintanawat et al found the odds of abuse among female elders who lived with family members, with a negative attitude towards elderly, were almost 10 times greater than elders living with family members with a positive attitude. In the same ways, the odds of being abused were 4.6 times greater among elders living with family members who were alcoholic or drug addicted, than who did not. The same as living in an inappropriate living status had the odds of falls three time higher than elders

who lived with a good living status(34). This research observed that family factors can result in violence but it did not specify that abuse causes falls. International literature reported that the consequences of elder abuse can result in falls(35) and the consequences are mostly contusion and in some cases severe injury(36). Falls caused by abandoned elderly, without necessary support for their physical activities such as walkers or glasses were concerned as one of types of elder abuse(37).

3.5 Distal Cause

3.5.1 Social and community support

Table 14 Descriptive data of social and community support of total 450 participants age >60 in the survey in Eastern of Thailand in 2018

Characteristic	N	%
Member of Senior Club	279*	62
Attending to Senior Club's activities	300*	66.7
Home-visit by Provincial administrative organization	52	11.6

*missing data=55

3.5.1.a Senior citizens club

Senior clubs in Thailand are introduced in the community-level with a reason of improving quality of life in the aging society, in which the club is aimed to organize activities, such as religious, cultural and health promotion(38). The result from the research survey showed 75 participants who were members of senior clubs fell (26.9%), this proportion was more than participants who were not members (n=29, 25%). Participants who attended senior club's activities fell more than those who did not (n=100, 28.2%). This study did not find a significant effect of senior clubs to falls among the elderly.

3.5.1.b Home visit by provincial Administrative Organizations

Among participants that were visited by the provincial administrative organization, 13 of them fell (25%) this proportion is lower than those who were not visited by provincial administrative organizations (n=108, 27.1%). This survey did not state which activities were done during the home visit, but according to the provincial administrative guideline had mentioned health promotion and health follow up activities, which might increase the health knowledge of the elderly. The analysis of this survey showed no association between home visiting by Provincial Administrative Organizations and falls.

While the large sample size study of Sophonratanapokin et al from the national survey, observed association in social-communities factors, that affect to falls among the elderly, but this study did not specify what kind of activity and where was it(19). I did not find the study explored home-visit by the community and its relation to falls.

3.5.2 Health system, Policy and Government factors

Table 15 Descriptive data the health insurance and support from government of total 450 participants age >60 in the survey in Eastern of Thailand in 2018

Characteristic	N	%
Health insurance		
- National Health Insurance	373	82.9
- Civil Health Insurance	58	12.9
- Security social Insurance	14	3.1
- Private Health insurance	5	1.1
Receive the elderly subsistence allowances	410	91.1
Home-visit by health facilities	173	38.4
Having the health recorded notebook	119	26.4

The analysis in the cause of falls in policy and governmental level is aimed to explore the effect of providing the elderly subsistence allowance (ESA), home-visiting by health facilities (HFs) and having the individual health recorded notebook, which provided from them. The Thai Government distributed ESA to the elderly monthly, some of them did not receive this due to various reasons. The result reported that participants who receive ESA, fell more (n=112, 27.3%) than those who did not receive (n=9, 22.5%). The same as participants who had been visited by HFs fell more (n=53, 30.6%) than those who had not been visited (24.5%). There was no report of association between receiving ESA, home visit by HFs and falls (see table 13).

The individual health record notebook is one of the health policies for the elderly, to record their health data, both at home and at the health facilities. Although this notebook is provided in health facilities for all elders, but not all of them keep and use it, this also depends on the concerning of individual health. The result showed 33 participants who had a health recorded notebook fell (27.7%), which proportion was slightly lower than participants who did not have. The notebook had also content of health education such as recommended exercise and nutrition, which will benefit elders to be healthier. No significant association between having a recorded book and falls was observed.

The result of this survey did not show the relation between health system, policy and health recorded notebook to falls and I did not find other research that is done in these issues to support the evidence of no association.

Table 16 Bivariate analysis of distal factors with falls in a survey in Eastern Thailand 2018

Characteristic	Falls (% ^a)	None-falls (% ^a)	OR	95% CI	p-value
Member of Senior Club				0.671-1.812	0.699
- Yes	75 (26.9)	204 (73.1)	1.103		
- No	29 (25)	87 (75)	1		
Attending to Senior Club's activities				0.080-2.364	0.146
- Yes	100 (28.2)	255 (71.8)	1.382		
- No	21 (22.1)	74 (77.9)	1		
Home-visit by Provincial administrative organization				0.460-1.741	0.444
- Yes	13 (25)	39 (75)	0.895		
- No	108 (27.1)	290 (72.9)	1		

Receive the elderly subsistence allowances				0.597-2.805	0.512
- Yes	112 (27.3)	298 (72.7)	1.295		
- No	9 (22.5)	31 (77.5)			
Home-visit by health facilities				0.889-2.073	0.157
- Yes	53 (30.6)	120 (69.4)	1.357		
- No	68 (24.5)	209 (75.5)	1		
Having the health recorded notebook				0.663-1.695	0.809
- Yes	33 (27.7)	86 (72.3)	1.060		
- No	88 (26.6)	243 (73.4)	1		

^a % of sub-group in each characteristics (in the row)

^b p-value<0.05: significant association

^c Result was analyzed by using Fisher's Exact Test

CHAPTER FOUR : FALLS PREVENTION AND BEST PRACTICES

In this chapter, I will review preventive intervention of falls in Thailand and other countries in different levels. Interventions that have purposes to prevent falls among elderly but did not evaluate the rate of falls at the result, were excluded in this chapter.

4.1 Prevention at the individual level

4.1.1 Motor and balance problems

The research conducted in Thailand, was aimed to find the effectiveness of simple balancing exercises in elderly outpatients, who had experienced falling in the previous year. Kuptniratsaikul et al had introduced participants to seven balancing exercises, which focused on improving strengthening, hip muscle, quadricep muscle, marching, stepping over a bench, standing from the chair and walking heel-to-toe. Participants were evaluated in a balancing ability at baseline, three, six, nine, and 12 months by physical status indicators. The final result of this study showed that 49% of participants did not fall after one year, which explained that the number of falls cases reduced in a one-year period. Likewise, the balancing abilities, which were secondary outcomes of this study, were significantly increased after intervention, except functional reach. This study had a limitation in the lacking of a control group to compare the intervention and participants were recruited from an advertising board in the hospital which might be a selection bias(39).

The randomized controlled trail study in New Zealand conducted the exercise intervention in 240 participants of the age 75 and above. This study had divided participants into exercise and control group. Exercises were muscle strengthening training and an improving balance program, which was introduced by trained nurses, participants were expected to do 30 minutes exercise three times a week. The assessment was done after one year, the result showed in the exercise group had a 46% decrease in number of falls comparing to the control group (IRR=0.54, 95%CI=0.32-0.90. This study explored the cost effectiveness which resulted explained that the exercise program among participants age 80 and older was cost saving \$NZ576 per fall case and \$NZ1563 per injurious falls case(40).

4.1.2 Correct visual impairment by Cataract surgery

Cataract surgery was introduced as preventive intervention of falls among the elderly in Vietnam. This cohort study (n=413) was conducted in cataract patients age above 50 and excluded participants who had previous surgery for cataract, cataract from diabetes, glaucoma or other ocular disease, and neurological impairment. The assessments were done to find the outcome, which is an incidence of falls before 1-3 months and in the year after surgery. The result was reported, reducing the prevalence of falls after 1 year in first-eye surgery 1.9% and 6.1% in both eyes surgery. After adjusted potential confounding, the risk of falls after one year decreased by 78% in first-eye surgery and 83% in both-eye surgery. The study did not indicate which variable they used to adjust confounding(41).

4.2 Prevention at the proximal level

The randomized controlled trail study (n=391) in southern Germany, introduced the modification of environmental hazards among frail elders after a one-year intervention. The program was implemented after the participant were discharged from the hospital with

different medical conditions. This program included regular home-visits in order to access the risks and to advise on possible modifications of the environmental hazard in the house. After a one-year assessment, the result showed a lower 31% of fall rate among the intervention group than the control group. (IRR=0.69, 95%CI=0.51-0.95). This study also illustrated that among participants who fell 2 time or more in the intervention group they had a 37% lower falls rate than in control group (IRR=0.64, 95%CI=0.37-0.99, p=0.047). While the group of participants who did not modify their house and control group explained no difference in the rate of falls. The highest percentage of compliance in home modification was a shower seat, following by an emergency call, grab bars, night light and anti-slipping mat in the bathroom (42).

4.3 Prevention at the distal level

This study explored intervention in distal level as the preventive program that was introduced in communities, care-facilities and hospitals. One intervention was introduced in the community-based level in Thailand (n=28) by Kittipimpanon et al. It was a combine fall prevention model which consisted of a falls risk assessment, a campaign, education, training for balance improvement, home visit for home hazardous modification and a fall management system. The intervention was implemented in the senior club of the communities and assessed a number of falls before and after a 10 month-period, which result reported a 24.86% reduction of falls incidence with an improvement of legs muscles strengthening, and balancing abilities(43). Although the study had small numbers of participants, it is aimed to find a long-term effect compared to other literature. The result after 3 years found an unchanged percentage of falls. Beside this study conducted qualitative research from program leaders and providers in order to evaluate the process of intervention. The result explained exercise was most effective long-term activity and factors of successful program were well-cooperation between the community, the health care worker in community level and dwellers, funding and recognition of falls prevention's importance(44).

To explore the intervention in care-facilities and hospitals, the Cochrane review studied among 95 randomized controlled trails (138164 participants age >65) and aimed to find a reduction in rates of fall and risk of falls from preventive interventions that were conducted in care facilities and hospitals in 23 countries. This study explored single interventions; exercise training, medication review, vitamin D supplementation, environment technology, social environment, education, psychological training, and multiple/multifactorial interventions. Among the intervention in care facilities, vitamin D supplementation had an effect on the falls reduction (RaR= 0.72, 95% CI=0.55-0.95), while other interventions reported no reduction of the falls rate with uncertainty in results due to inadequate evidence quality. Among interventions in hospitals, additional exercise with physiotherapy could result in a decreasing risk of falling, although it was reported uncertain due to a very low quality of evidence(45).

CHAPTER FIVE : DISCUSSION AND LIMITATION

5.1 Discussion

5.1.1 General

Representativeness of the sample size, comparison with the national level and context

This exploring study covered the Eastern region of Thailand, this area has special characteristic in manufacture industries growing with the increase in the working-age group. This situation brought the Eastern region a different proportion of age-groups from other regions and the percentage of older population was reported the second lowest in the national report (14.85%) in 2019(7). The ratio of male to female elders were approximately 3:4 (7), while the survey had a different proportion (1:4). Few male respondents were explained by the survey researchers that most of them went out for work when the survey was conducted. The same reason can be explained in lower percentage of respondents in age group 60-69 and higher percentage of age group 80 and above comparing to the regional statistic data.

Falls in the survey

The rate of falls among the elderly in the survey was 26.9 %, which is higher than studies in the urban community in Bangkok(19.8%) and in a Southern province(26.1%)(46)(20). The incidence of falls worldwide is 28-35% and this depends on each country such as Chile is 34% while Japan has 20% (47). The elderly fell more inside the house and the most common place is on the balcony. Consequences of falls were in majority injuries, while only a few cases had bone fractures and head laceration and wounds. Half of the elderly did not go to health care facilities and self-cured.

5.1.2 Individual level

Dependence factors

Although this study did not find a significant association between age and falls in the survey, the higher rate of falls was reported in the older age group. Thai literatures explained that older age had a higher chance of falls with a strong association(48), due to degenerative changes in physical, sensory and cognitive functions including more possibilities of having medical conditions(49).

A number of Thai articles reported an association in that the female gender had a higher chance of falls as same as international literatures (48)(50)(51), while the result of secondary data analysis did not find an association in the survey, even though female elders fell more than males. Different genders have different physical activities, lifestyle and circumstance which influences risks of falls differently(50). Although women had higher incidences of falls, men seem to have more severity, due to greater hospitalization and a mortality rate than women(52).

A study of other demographic status (education, marital status, religion, occupation and income) in the survey was not reported in association to falls, compared with a number

of literatures in Thailand found low levels of education resulted in a higher prevalence of falls in older adults. A higher level of education can lead to a reduced chance of falls due to less exposure to harmful jobs and a better health education and health status(20). Insufficient income was reported in relation to an increasing risk of falls in the Thai study. The same result was found in the Mexican American study (53) in which influences were poor environment, poor nutrition intake and access to health care services(52).

According to the data analysis of physical status and medical conditions, the result found evidence of significant association of BMI, excessive waist circumference, balancing impairment. Hearing impairment can result in falls in the survey as same as the result of a study in the United Arab Emirates and Taiwan(54)(55). Sleeping pills are considered in the group of antidepressant medication, which have been a concern as a risk factor of fall according to the survey as same as an international literature(55). The reason of their side effect causing postural hypotension and some pharmacological groups can cause decreasing bone mineral density, which can lead to falls(56).

Other literatures in Thailand found, visual impairment, OA knee, foot disease strongly associated to falls. Visual impairment can lead to falls among elders directly(57) and indirect influence by decreasing mobility and balancing ability, which finally had an impact on falls(58). The result in the Thai literature review explained neurological disorders such as stroke, Parkinson's disease and dementia are risk factors of falls, likewise the similarly result of Finkelstein et all's study explained the decline of the cognitive function can result in falls among elderly patients which further observed rising in cost of consequence treatment(59).

Independence factors

Although healthy lifestyle was not reported in association to falls in the secondary data analysis of the survey, there was evidence that doing regularly exercise can reduce the incidence of falls in the Thai article. Elders who did not exercise regularly tend to fall more, the assessment of their physical performance were lower than the non-falls elders(55). In the same way, exercise can reduce the chance of falls in terms of controlling BMI, waist circumference, improving balance, muscle and joint function(60). Another evidence explained the benefit of exercise to falls, that it has an effect on increasing bone density in postmenopausal older women(61).

In this study I did not find that a level of HL has effect to falls, this is the first study to evaluate this association in Thailand, there is no other evidence to support this conclusion. Fear of falls described the feeling that many older adults experienced, due to their history of falls, co-morbid conditions and functional impairments (62). Similarly, the study in Canada found that the elders fear of falls can result in a decrease in physical activities, muscle weakness and finally lead to falls(57). Nevertheless, this can be positive factor when the elderly concern in falls and consequences, then they attempt more to prevent it (52).

5.1.3 Proximal level

Environmental hazardous

Environmental factors have a major impact on falls among the elderly in this study the same as in other international studies(63)(32). A hazardous Indoor home is crucial, as many studies had reported elders fell more inside the house. This study is aimed to explored the

risk of falls in different type of the houses, including a raised floor house, that contains stairs and steps, which might be harmful for the elderly. Although data analysis of the survey did not show that the characteristic of the house has an impact on falls, the literature review in hip fracture patients found that multi-stories floor are a risk factor especially for the severe cases. Other literatures in Thailand emphasized more on rooms, types of the floor, and handrails. Kitchen, bathroom, bedroom are places that have a high risk of falls in the house. Same as a slippery floor and a slope way, which characteristics can lead to losing balance control and end up with falls(31)(32)(63). Having a handrail in the bathroom is a protective factor of falls according to literature review, when elders who have bathroom without this support they seem to fall more(32)(64). Literature review found dim light inside the house is one of the factors related to falls, which can lead to reduced visual ability of the elderly and might result in falls at the end.

The character around the house with rough ground with rocks, unclear area, with discarded items and with an incline pathway are risk factors of falls among older adults according to the survey. These home hazards are essential, while the incidence of falls can be increased when two or more risks are combined(63).

Family support and violence from family members

Although, the survey did not find the relation of family support, the living condition has an impact on falls, when elders who live alone tend to fall more than those who live with their spouses, according to Thai literature. Similarly result explained that living alone can increase the risk of falls, which result might depend on types of house(65), on the other hand, living with spouses can reduce the incidence of falls(66).

Living with a family might have a negative effect on the elderly because it can result in abuse from family members. A high prevalence of elder abuse was reported in Thailand, it becomes a common problem in many countries and most of the cases are caused by family members(67). Negative thoughts of family members play a big role in violence among the elderly, besides, elders who have poor living conditions or live with alcoholic, drug addict family members have a higher chance of abuse. Although living with family member can result in violence there is no well-defined evidence stated that elders abuse caused by family member has the outcome of falls.

5.1.4 Distal level

Social and community factors

Senior clubs provide activities in the community level that are aimed to improving life and health status among elders, through health education and other activities. The engagement of senior club activities in the Eastern region did not relate to falls in the survey and there is no literature exploring the effect of this factor to falls. The same as home-visiting by the community is not related to falls, but elders who had been home-visited by communities seems to falls less, which might be explained by health promotion and health follow up activity during home-visits, can increase awareness and reduce the fall rate. This may have a selection bias when falls rate was low in specific group of participants.

Government and policy factors

This study of data analysis explored that receiving ESA, home visit by HF and having a health recorded notebook, which was introduced by the ageing health policy, did not have an effect on falls among the elderly. Even though, there was lower rate of falls among those who used the health recorded notebook, which might be related to concerns of their health, which leads to more prevention in falls than others. These implementations are crucial in the ageing field; no study reconnoitred its effect to falls. Home-visits by HF were mostly reported as a part of the falls prevention programs, that have an impact on the reducing falls prevalence (68)(42).

5.1.5 Preventive interventions of falls

Exercise was found an effective intervention to improve muscle strength and balance ability of the elderly and reduce the rate of falls among elders. Exercise is found to be a cost-saving intervention of falls among the elderly in developed countries, especially in elders aged above 80. Thailand, middle-income country may have different results due to a different financial status and no study is done to explore this. Cataract surgery also benefits in falls prevention because this will improve visual ability of elders(69). Interventions in the environmental factors are included in both risk assessment and modification hazardous homes are found to be efficient. Most of the home modifications are removing the hazardous situation, install handrails in the toilet and the stair railing and set the call for a help system for an emergency situation(47).

A combined preventive program in the community level is effective in a long-term reduced number of falls. Especially exercise training is a concern as continuous activity among elders, while a home hazardous modification was found limiting, due to its cost. To prevent falls in care-facilities and hospitals, vitamin D supplementation results in a falls rate reduction and additional exercise combined with physiotherapy can reduce the risk of falls among the elderly. These factors have an unclear effect on falls prevention due to the poor quality of the evidences.

5.2 Limitations of the study

The survey is a study to cover all provinces in the eastern part including 1 province in the central part, according to the health region in Thailand. The sampling province, which were included in the survey, were only provinces in the eastern area. The research survey included only older participants who were literate and physically active. The proportion of male and female respondents were different from regional ratio due to many male elders were absent during data collection.

The survey was aimed to explore factors related to a healthy lifestyle among the elderly, which included the data of falls, but some of the data was not specific. There were missing data in membership of senior club and attendance to senior club's activity, which were excluded during the analysis. In addition to this, the data of the history of falls was collected within 1 year, which may be a recall bias, due to limited memorial abilities of the elderly. Data of insufficient income, medical condition, physical status, and hazardous environments were collected by what participants answer. There was no observation at houses or collection data from a medical record which might be measurement bias in the result of the survey. The data in the HL level was not covered, all 6 categories of health literacy guideline itself and nowadays there was no guideline tool to assess the HL level. This

survey is limited to analyze with adjusting confounding due to the small sampling size of data, several literatures in Thailand were analyzed and explained about this.

I did not find Thai literature that explored the impact of senior clubs and using the elderly recorded notebook on falls, as same as elder abuse. Few studies in elderly abuse in Thailand were published, but all of them were not study in falls consequences and in articles concerned events were caused by family members.

The causation conceptual framework was useful to find factors in each level and the linkage between. I have found that the health system, government and policy factor in distal level can link to the social-community factor as same as it can link to independent individual and environmental factors.

CHAPTER SIX : CONCLUSION AND RECOMMENDATION

6.1 Conclusion

Falls are an unintended injury that happens more often among the elderly due to degenerative changes, both physically and mentally. Becoming an ageing society in Thailand, falls incidence, is high and continuing to increase, which problem is concerned in Thai MoPH. Especially in the eastern region of Thailand, where the social and financial context is slightly different from other regions, due to the growing of the manufacturing industry that affects families and living conditions. The incidence of falls in the East of Thailand is higher than other regions, with limited study in ageing health and related factors to falls that could help to improve this problem.

The individual cause of falls is beyond each of elders themselves. Age is one of the risk factors of falls, when the older age group falls easier than younger, due to physical and functional changes and other co-morbid conditions that are increasing with older age. Female gender affects the falls incidence more, because of higher proportion comparing to males and females' activities, lifestyle and circumstances could lead to falls more than male. In addition to this, with a longer life expectancy in the female gender this bring more females in the old age group, which is an obviously rise up of the incidence of falls among this gender. Males fall less but the consequences seem to be more severe than female. Most of the demographic status namely, marital status, religion, occupation are not related to falls, while low education level and insufficient income are risks factors of fall, due to this increases the exposing to dangerous work, and poor life quality.

Physical status plays a big role in falls, when elders with a higher BMI or excessive waist circumference are more likely to falls than who have a normal level. Impairment of visual and hearing function are factors that lead to falls among the elderly. Sleeping pills are concerned as a risk factor of falls, when these are more prescribed to older adults. Several medical conditions that affect the reducing of the physical function such as foot disorder, knee OA, can lead to falls among the elderly. In the same way, the decreasing of neurological function, such as Alzheimer's disease, dementia, Parkinson's disease and stroke, have an effect on the increasing chance of falls. A healthy lifestyle is a protective factor of falls, because eating vegetables regularly and not drinking alcohol can reduce falls rates. Doing exercise regularly will help the elderly to improve their muscle strength, balance and physical ability, which will decrease the risk of falls through improving the bone mass mechanism and this is also reducing the chance of osteoporosis.

Home hazard is a risk factor to falls and the common cause of falls in the house are slippery floors, steps, stairs, slope way, none-handrails or supporters, poor lighting. While many elders fall outside the house, because of unclear areas, rough ground with rocks and areas with discarded items or inclined pathways. Many preventive interventions aim to assess the risk and modify the house, due to high efficiency, however there is limited study that shows cost effectiveness in this intervention.

Living alone has a higher chance of falls than living with spouses or their children due to the care or support from them. Although staying with family members can be a protective factor of falls, this can also lead to elder abuse, which might result in falls. While limited evidence showed engaging to social activities is related to falls, many interventions aim to introduce in the community-based level such as exercise training, home hazardous assessment, and the health promotion program. The same as factors of falls in the policy and

governmental level are unknown because of limited research had been explored though this level.

6.2 Recommendation

This study recommends to develop in ageing health and will be described recommendations in different level of responsibilities.

Recommendation for Thai MoPH:

- Develop falls intervention in policy level. Policy option of falls prevention should be different due to different risks of falls between different age group and genders.
- Introduce calcium and vitamin D in the elderly diet guideline, to prevent falls.
- Introduce guideline of falls prevention including environmental hazards modification in HFs, care-facilities and in communities.

Recommendation for health facilities and health care workers and community:

- Cautioning doctors to prescribe sleeping pills, in order to avoid its side effect.
- Provide a health education that will increase the knowledge and understanding to elders and also their family members or care givers.
- Introduce combination program of intervention in community-level, which include environmental hazardous assessment and modification, health promotion, and exercise. However, excessive exercise can also cause falls, which need to be explained to the elderly to be aware.

Research recommendation:

- Research in the HL level and elder abuse to find the descriptive data, relevant cause and effect to health status including falls.
- Research in community and policy level both qualitative and quantitative in senior club's activities home-visit by HF and communities, having the health recorded notebook and other ageing health strategies. In order to find their relation to health status including falls, effectiveness, strength and weakness.
- Research in effectiveness of preventive intervention in exercise training, home hazardous modification including prevention in policy level.

References:

1. United Nations Statistics Division. UN data: Country Profile: Thailand [Internet]. UNdata. 2020 [cited 2020 Aug 6]. Available from: <http://data.un.org/en/iso/th.html>
2. Office of Royal Society. Geographic regions in Thailand [Internet]. 2015. Available from: <http://www.royin.go.th/?knowledges=การแบ่งภูมิภาคทางภูมิศาสตร์>
3. The World Bank Group. Thailand Overview [Internet]. 2019 [cited 2020 Aug 6]. Available from: <https://www.worldbank.org/en/country/thailand/overview>
4. World Bank Group. GINI index (World Bank estimate) - Thailand | Data [Internet]. 2019. [cited 2020 Aug 9]. Available from: <https://data.worldbank.org/indicator/SI.POV.GINI?end=2018&locations=TH&start=1981>
5. Office of the National Economic and Social Development Council. The Fifth National Economic and Social Development Plan [Internet]. 2019 [cited 2020 Aug 6]. Available from: https://www.nesdb.go.th/nesdb_en/ewt_w3c/ewt_dl_link.php?filename=develop_issue&nid=3780
6. National Statistic Office. Demography population and housing branch [Internet]. 2019 [cited 2020 Aug 9]. Available from: <http://statbbi.nso.go.th/staticreport/page/sector/th/01.aspx>
7. Department of Older Persons. Statistics of the Thai elderly 2019 [Internet]. 2019 [cited 2020 Jul 22]. Available from: <http://www.dop.go.th/th/know/side/1/1/275>
8. Thai Mistry of Public Health. Public Health Statistic [Internet]. 2018 [cited 2020 Aug 6]. Available from: https://bps.moph.go.th/new_bps/sites/default/files/statistic_61.pdf
9. World Health Organization. Mortality Report Thailand [Internet]. 2016 [cited 2020 Aug 9]. Available from: https://www.who.int/nmh/countries/tha_en.pdf?ua=1
10. The World Bank. Thailand: Sustaining Health Protection for All [Internet]. 2012 [cited 2020 Aug 6]. Available from: <https://web.archive.org/web/20120903000731/http://www.worldbank.org/en/news/2012/08/20/thailand-sustaining-health-protection-for-all>
11. World Health Organization. Preface Overview Humanity's Aging Living Longer New Disease Patterns Longer Lives and Disability New Data on Aging and Health Assessing the Cost of Aging and Health Care Changing Role of the Family Suggested Resources. Global Health and Aging. 2010.
12. WHO. Ageing and health [Internet]. 2018 [cited 2020 Feb 10]. Available from: <https://www.who.int/news-room/fact-sheets/detail/ageing-and-health>
13. Foundation of Thai Gerontology Research and Development Institute. Situation of the Thai elderly 2016 [Internet]. 2016 [cited 2020 Jan 29]. Available from: http://www.dop.go.th/download/knowledge/th1512367202-108_0.pdf
14. Pignolo RJ, Mary ·, Forcica A, Johnson JC, Ahn J. Aging Medicine Series Editors: Fractures in the Elderly A Guide to Practical Management Second Edition [Internet]. 2018 [cited 2020 Jun 11]. Available from: <http://www.springer.com/series/7622>
15. National Health Service. Falls [Internet]. 2018 [cited 2020 Jan 28]. Available from: <https://www.nhs.uk/conditions/falls/>
16. ThaiNCD. The statistic data of falls among elderly [Internet]. 2017 [cited 2020 Jan 29]. Available from: <http://www.thaincd.com/2016/news/hot-news->

- detail.php?id=12925&gid=18
17. Health System Research Institute (HRSI). Falls in Elderly [Internet]. 2014 [cited 2020 Jan 12]. Available from: <https://www.hsri.or.th/people/media/care/detail/5532>
 18. Murray CJL, Ezzati M, Lopez AD, Rodgers A, Vander Hoorn S. Comparative quantification of health risks: Conceptual framework and methodological issues. Vol. 1, Population Health Metrics. BioMed Central; 2003. p. 1.
 19. Sophonratanapokin B, Sawangdee Y, Soonthorndhada K. EFFECT OF THE LIVING ENVIRONMENT ON FALLS AMONG THE ELDERLY IN THAILAND. Vol. 43. 2012.
 20. Worapanwisit T, Prabpai S, Rosenberg E. Correlates of Falls among Community-Dwelling Elderly in Thailand. *J Aging Res.* 2018;
 21. Maneeprom N, Taneepanichskul S, Panza A. Fall among physically active elderly in senior housings, Bangkok, Thailand: situations and perceptions. *Clin Interv Aging* [Internet]. 2018 Oct 26 [cited 2020 Feb 24]; Volume 13:2149–59. Available from: <https://www.dovepress.com/fall-among-physically-active-elderly-in-senior-housings-bangkok-thaila-peer-reviewed-article-CIA>
 22. Chaiwanichsiri D, Janchai S, Tantisiriwat N. Clinical Section Foot Disorders and Falls in Older Persons. *Gerontology* [Internet]. 2009 [cited 2020 Feb 24]; Available from: www.karger.com
 23. Thaweewannakij T, Suwannarat P, Mato L, Amatachaya S. Functional ability and health status of community-dwelling late age elderly people with and without a history of falls. *Hong Kong Physiother J* [Internet]. 2016 Jun 1 [cited 2020 Jul 8];34:1–9. Available from: <http://dx.doi.org/10.1016/j.hkpj.2015.08.0011013-7025/BY-NC-NDlicense>
 24. Kuhirunyaratn P, Jindawong B. FACTORS RELATED TO FALLS AMONG COMMUNITY DWELLING ELDERLY. Vol. 44. 2013.
 25. Muangpaisan W, Suwanpatoomlerd S, Srinonprasert V, Sutipornpalangkul W, Wongprikron A, Assantchai P, et al. Causes and Course of Falls Resulting in Hip Fracture among Elderly Thai Patients. Vol. 98, *J Med Assoc Thai.* 2015.
 26. KPI MoPH. Percentage of healthy lifestyle of the elderly [Internet]. KPI webpage. 2020 [cited 2020 Jun 29]. Available from: <http://healthkpi.moph.go.th/kpi/kpi-list/view/?id=1492>
 27. Thai Health Promotion Foundation. Reduce belly-Reduce disease Guideline. 2014.
 28. Folk Doctor Foundation. Healthy Elderly with Food, Exercise and Emotion [Internet]. 2016 [cited 2020 Jul 21]. Available from: www.doctor.or.th
 29. Friedman SM, Munoz B, West SK, Rubin GS, Fried LP. Falls and fear of falling: Which comes first? A longitudinal prediction model suggests strategies for primary and secondary prevention. *J Am Geriatr Soc* [Internet]. 2002 Aug [cited 2020 Aug 4];50(8):1329–35. Available from: <https://pubmed.ncbi.nlm.nih.gov/12164987/>
 30. Thiamwong L, Suwanno J. Fear of Falling and Related Factors in a Community-based Study of People 60 Years and Older in Thailand. *Int J Gerontol.* 2017 Jun 1;11(2):80–4.
 31. Yu PL, Qin ZH, Shi J, Zhang J, Xin MZ, Wu ZL, et al. Prevalence and related factors of falls among the elderly in an urban community of Beijing. *Biomed Environ Sci* [Internet]. 2009 Jun [cited 2020 Jul 29];22(3):179–87. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S089539880960043X>
 32. Cesari M, Landi F, Torre S, Onder G, Lattanzio F, Bernabei R. Prevalence and Risk Factors for Falls in an Older Community-Dwelling Population [Internet]. Vol. 57, *Journal of Gerontology.* 2002 [cited 2020 Jul 29]. Available from:

- <https://academic.oup.com/biomedgerontology/article-abstract/57/11/M722/625753>
33. Kulachai W. Elder Abuse in Thailand: Prevalence, Causes, Consequences, and Public Supports. *Int J Manag Appl Sci* [Internet]. 2018 [cited 2020 Jul 27];4(10):52–26. Available from: https://www.researchgate.net/publication/328095961_Elder_Abuse_in_Thailand_Prvalence_Causes_Consequences_and_Public_Supports
 34. Chintanawat R, Sucamvang K, Sirirat Panuthai S. Prevalence and Risk Factors for Abuse among Thai Female Older Adults. *Thai J Nurs* [Internet]. 2006 [cited 2020 Jul 27];21(31–46). Available from: <https://he02.tci-thaijo.org/index.php/TJONC/article/view/2390/2290>
 35. Kavak RP, Meltem Özdemir . Radiological appearance of physical elder abuse. 2019 [cited 2020 Jul 9];10:871–8. Available from: <https://doi.org/10.1007/s41999-019-00246-5>
 36. Cham GWM, Seow E, Cham G. The Pattern of Elderly Abuse Presenting to an Emergency Department. Vol. 41, *Singapore Med J*. 2000.
 37. Gibbs LM. Understanding the Medical Markers of Elder Abuse and Neglect: Physical Examination Findings. [cited 2020 Jul 10]; Available from: <http://dx.doi.org/10.1016/j.cger.2014.08.002>
 38. Senoir citizen council of Thailand. Guideline of senior citizen club [Internet]. 2016 [cited 2020 Jul 17]. Available from: http://www.dop.go.th/download/knowledge/knowledge_th_20161706104732_1.pdf
 39. Kuptniratsaikul V, Praditsuwan R, Assantachai P, Ployetch T, Udompunturak S, Pooliam J. Effectiveness of simple balancing training program in elderly patients with history of frequent falls. *Clin Interv Aging*. 2011;6(1):111–7.
 40. Robertson MC, Devlin N. Effectiveness and economic evaluation of a nurse delivered home exercise programme to prevent falls. 1: Randomised Controlled Trial - in - *BMJ : British Medical Journal (Clinical Research edition)*. *BMJ Br Med J (Clinical Res Ed)* [Internet]. 2001 Mar 24 [cited 2020 Aug 6];322(7288):697. Available from: [http://atoz.ebsco.com/Titles/SearchResults/12084?IsFromAdvancedSearch=True&Find=BMJ : British Medical Journal \(Clinical Research edition\)&GetResourcesBy=TitleNameSearch&resourceTypeName=allTitles&resourceType=&SearchType=ExactMatch](http://atoz.ebsco.com/Titles/SearchResults/12084?IsFromAdvancedSearch=True&Find=BMJ : British Medical Journal (Clinical Research edition)&GetResourcesBy=TitleNameSearch&resourceTypeName=allTitles&resourceType=&SearchType=ExactMatch)
 41. To KG, Meuleners L, Bulsara M, Fraser ML, Van Duong D, Van Do D, et al. A longitudinal cohort study of the impact of first- and both-eye cataract surgery on falls and other injuries in Vietnam. *Clin Interv Aging* [Internet]. 2014 Apr 28 [cited 2020 Aug 6];9:743–51. Available from: </pmc/articles/PMC4011894/?report=abstract>
 42. Nikolaus T, Bach M. Preventing Falls in Community-Dwelling Frail Older People Using a Home Intervention Team (HIT): Results From the Randomized Falls-HIT Trial. *J Am Geriatr Soc* [Internet]. 2003 Mar 1 [cited 2020 Jul 13];51(3):300–5. Available from: <http://doi.wiley.com/10.1046/j.1532-5415.2003.51102.x>
 43. Kittipimpanon K, Amnatsatsue K, Kerdmongkol P, Maruo SJ, Nityasuddhi D. Development and Evaluation of a Community-based Fall Prevention Program for Elderly Thais. Vol. 16, *Pacific Rim Int J Nurs Res*.
 44. Kittipimpanon K, Kraithaworn P. The effectiveness of community-based fall prevention model to physical performance and falls among older adult in urban community Bangkok. *J Public Heal Nurs* [Internet]. 2015 [cited 2020 May 9];29:98–113. Available from:

- <https://med.mahidol.ac.th/nursing/sites/default/files/public/research/Nation/2557/FullTxt/fullTxtNo49.pdf>
45. Cameron ID, Dyer SM, Panagoda CE, Murray GR, Hill KD, Cumming RG, et al. Interventions for preventing falls in older people in care facilities and hospitals [Internet]. Vol. 2018, Cochrane Database of Systematic Reviews. John Wiley and Sons Ltd; 2018 [cited 2020 Aug 6]. Available from: </pmc/articles/PMC6148705/?report=abstract>
 46. Assantachai P, Praditsuwan R, Chatthanawaree W, Pisalsarakij D, Thamlikitkul V. Risk factors for falls in the Thai elderly in an urban community. *J Med Assoc Thai* [Internet]. 2003 Feb [cited 2020 Feb 11];86(2):124–30. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/12678149>
 47. World Health Organization. Ageing and life course, family and community health WHO Global report on falls prevention in older age. 2007.
 48. Chang VC, Do MT. Original Contribution Risk Factors for Falls Among Seniors: Implications of Gender. *Am J Epidemiol* [Internet]. 2015 [cited 2020 May 9];181(7):521–31. Available from: <https://academic.oup.com/aje/article-abstract/181/7/521/150122>
 49. Bueno-Cavanillas A, Padilla-Ruiz F, Jiménez-Moleón JJ, Peinado-Alonso CA, Gálvez-Vargas R. Risk Factors in Falls among the Elderly According to Extrinsic and Intrinsic Precipitating Causes. Vol. 16, *European Journal of Epidemiology*. 2000.
 50. Duckham RL, Procter-Gray E, Hannan MT, Leveille SG, Lipsitz LA, Li W. Sex differences in circumstances and consequences of outdoor and indoor falls in older adults in the MOBILIZE Boston cohort study [Internet]. 2013 [cited 2020 Jul 23]. Available from: <http://www.biomedcentral.com/1471-2318/13/133>
 51. Gale CR, Cooper C, Aihie Sayer A. Prevalence and risk factors for falls in older men and women: The English Longitudinal Study of Ageing. *Age Ageing*. 2016;45(6).
 52. World Health Organization. WHO Global report on falls Prevention in older Age. 2007.
 53. Reyes-Ortiz CA, Snih S Al, Loera J, Ray LA, Marides K. Risk factors for falling in older Mexican Americans. *Ethn Dis* [Internet]. 2004 [cited 2020 Jul 23];14. Available from: https://www.researchgate.net/publication/8383143_Risk_factors_for_falling_in_older_Mexican_Americans
 54. Sharif SI, Al-Harbi AB, Al-Shihabi AM, Al-Daour DS, Sharif RS. Falls in the elderly: Assessment of prevalence and risk factors. *Pharm Pract (Granada)*. 2018;16(3).
 55. Huang HC, Gau ML, Lin WC, Kernohan G. Assessing Risk of Falling in Older Adults. *Public Health Nurs*. 2003 Sep;20(5):399–411.
 56. Hill KD, Wee R. Psychotropic drug-induced falls in older people: A review of interventions aimed at reducing the problem [Internet]. Vol. 29, *Drugs and Aging*. 2012 [cited 2020 Jul 24]. p. 15–30. Available from: <http://link.springer.com/10.2165/11598420-000000000-00000>
 57. Public Health Agency of Canada. Seniors’ Falls in Canada (Second Report) [Internet]. 2014 [cited 2020 Jul 23]. Available from: <https://www.canada.ca/en/public-health/services/health-promotion/aging-seniors/publications/publications-general-public/seniors-falls-canada-second-report.html#s3-1>
 58. Ray CT, Wolf SL. Review of intrinsic factors related to fall risk in individuals with visual impairments. Vol. 45, *Journal of Rehabilitation Research and Development*. 2008. p. 1117–24.
 59. Finkelstein E, Prabhu M, Chen H. Increased prevalence of falls among elderly

- individuals with mental health and substance abuse conditions. *Am J Geriatr Psychiatry*. 2007 Jul 1;15(7):611–9.
60. Gardner MM, Robertson MC, Campbell AJ. Exercise in preventing falls and fall related injuries in older people: A review of randomised controlled trials [Internet]. Vol. 34, *British Journal of Sports Medicine*. *Br J Sports Med*; 2000 [cited 2020 Jul 24]. p. 7–17. Available from: <https://pubmed.ncbi.nlm.nih.gov/10690444/>
 61. Day L, Fildes B, Gordon I, Fitzharris M, Flamer H, Lord S. Randomised factorial trial of falls prevention among older people living in their own homes. *Br Med J* [Internet]. 2002 Jul 20 [cited 2020 Jul 24];325(7356):128–31. Available from: </pmc/articles/PMC117228/?report=abstract>
 62. Rachel M, Birks Y, Hall H, Torgerson D, Watt I. Exploring the relationship between fear of falling and neuroticism. *Age Ageing* [Internet]. 2006 [cited 2020 Jul 24];35(2):143–7. Available from: <https://academic.oup.com/ageing/article-abstract/35/2/143/27936>
 63. Rubenstein LZ. CLINICAL RISK ASSESSMENT, INTERVENTIONS AND SERVICES Falls in older people: epidemiology, risk factors and strategies for prevention. *Age Ageing* [Internet]. 2006 [cited 2020 Jul 24];37–41. Available from: https://academic.oup.com/ageing/article-abstract/35/suppl_2/ii37/15775
 64. CDC. Preventing Falls: How to Develop Community-based Fall Prevention Programs for Older Adults. 2015.
 65. Todd C, Skelton D. What are the main risk factors for falls amongst older people and what are the most effective interventions to prevent these falls? [Internet]. 2004 [cited 2020 Jul 30]. Available from: <http://www.euro.who.int/document/E82552.pdf>,
 66. Van Bommel T, Vandenbroucke JP, Westendorp RGJ, Gussekloo J. In an observational study elderly patients had an increased risk of falling due to home hazards. *J Clin Epidemiol* [Internet]. 2005 Jan 1 [cited 2020 Jul 27];58(1):63–7. Available from: <http://www.jclinepi.com/article/S0895435604002276/fulltext>
 67. Giurani F, Hasan M. Abuse in elderly people: The Granny Battering revisited. *Arch Gerontol Geriatr*. 2000 Dec 1;31(3):215–20.
 68. Huang T-T, Acton GJ. Effectiveness of Home Visit Falls Prevention Strategy for Taiwanese Community-Dwelling Elders: Randomized Trial. *Public Health Nurs* [Internet]. 2004 May 1 [cited 2020 Jul 27];21(3):247–56. Available from: <http://doi.wiley.com/10.1111/j.0737-1209.2004.21307.x>
 69. Larson L, Bergmann TF. Taking on the fall: The etiology and prevention of falls in the elderly [Internet]. Vol. 11, *Clinical Chiropractic*. 2008 [cited 2020 Jul 29]. p. 148–54. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1479235408000382>

Acknowledgement

I would like thanks to the owner of the survey for my secondary data analysis part in my thesis and thanks to Health Center 6, the organization of the research owner.

I would like to thank to my thesis advisors for all supports with the thesis process. I am thankful to my academic advisor, KIT staff, KIT and Nuffic that support this course.

I am grateful to have international colleagues in ICHD course, giving advice, working together, sharing experiences which helps me to develop skill and knowledge.

Annex 3

Permission letter to use research survey data

To: Ploi Kongkoot

Royal Tropical Institute,
Mauritskade 64, Amsterdam,
The Netherlands

Date: 27 May 2020


Dear Ploi,

REPLY TO THE PERMISSION LETTER TO USE RESEARCH SURVEY DATA

Thank you for your interest in the data from my research survey "The related factors to health status and behavior in elderly in the Eastern part of Thailand". I appreciate that you find it useful for further study in aging health in the Thai population and I am glad to permit you to use my data for your study. Your research topic is interesting and I hope that after all, it will benefit the elderly health in Thailand.

I sincerely hope that your attempt will meet with great success. If you need any clarification or advice, please feel free to contact me.

Yours sincerely,


Tippawan Jumpang
Email: kaeja2521@hotmail.com

Waiver letter

From Ploi Kongkoot
Royal Tropical Institute
Mauritskade 64, Amsterdam,
The Netherlands

11 June 2020,

To: Chair Research Ethics Committee, KIT

Dear Pam Baatsen ,

This letter is to request a waiver of ethical clearance for a study on Factors related to Falls among the Elderly in the Eastern, Thailand. The data used is derived from the research survey "The related factors to health status and behavior in elderly in the Eastern part of Thailand".

The study is done in 2018 by Tippawan Jumpang, registered nurse professional level at Health Center 6, Thailand. The research survey was done with the purpose of exploring health status and healthy lifestyle of the elderly in the health region 6 (Eastern part of Thailand) and to find the related factors according to these issues. A secondary data analysis will be done by Ploi Kongkoot, in the context of KIT thesis research. The purpose of the study is to explore the related factor to falls among the elderly in the Eastern region of Thailand. The study focuses on individual factors, environmental factors, Family-social-community support factors, which cause falls in the old age group population. This secondary data analysis will be emphasized more on the outcome of falls among elderly comparing with the research survey that focuses on general health status and healthy lifestyle. The study results will be used to improve the intervention to reduce the incidence and prevalence of falls among the elderly in Thailand.

I would like to kindly request the Research Ethical Committee to a waiver of ethical clearance for this study for the following reasons:

1. The data has been anonymized.
2. Data (use) has been approved by the data owner, Tippawan Jumpang.
3. Informed consents were given and it would be feasible or practicable to ask informed consent to the participants to whom the data belong.
4. The research has important social, educational, and/or scientific value, namely exploring all the factors related to falls among elderly and best practices in the prevention of falls. The result of the study will be useful for the further plan of intervention and giving the recommendation to the Thai Ministry of Health in order to reduce the incidence of falls in the future.
5. The research poses no more than minimal risks to participants, including social and cultural risks such as discrimination, stigmatization or psychological discomfort or harm; and does not give rise to the disclosure of the participant's identity for privacy. The collected data from participants will be ensured securely keeping. The research

survey's data were asked the permission to use for this study and the data will proceed only after receiving the agreement from the owner of the research. The data can be accessed only authorized person and will only be used for the thesis research purpose.

I hope to have informed you sufficiently on the objective and content of this study to make a decision on my request.

Yours sincerely,

Ploi Kongkoot

Ploi Kongkoot

Co-Signer of academic advisor : Lisanne Gerstel

Co-Signer Signature

L Gerstel

Date: 11 June ,2020

Clearance letter for waiver



RESEARCH ETHICS COMMITTEE

Contact: Meta Willems (secretary REC)
Telephone +31 (0)20 568 8514
m.willems@kit.nl

To: Ploi Kongkoot
KIT ICHD Student
By E-mail: pl_oy_2k@hotmail.com

Amsterdam, 16 June 2020

Subject Decision Research Ethics Committee regarding a waiver for a study on "Factors related to falls among the elderly in Eastern region in Thailand" (S-126)"

Dear Ploi Kongkoot,

The Research Ethics of the Royal Tropical Institute (REC) has reviewed your application for a waiver for a study on "Factors related to falls among the elderly in Eastern region in Thailand" (S-126)" that was submitted on 9 June, 2020.

Your proposal has been exempted from full ethical review based on the following considerations:

1. the data has been anonymized and will be stored in a safe place, only accessible for you for the purpose of your thesis;
2. the data has been approved by the data owner being Tippawan Jumpang a registered nurse at the professional level, working for Health Center 6, the Department of Health in Thailand, who conducted the original study in 2018;
3. informed consent was obtained from the participants for the original study, while the current study objectives are aligned with the objectives of that original study.
4. the analysis has an important public health function;
5. the research poses no more than minimal risks to participants and does not give any rise to the disclosure of the participant's identity.

The Committee grants this waiver provided that you inform the KIT GDPR project officer about your research for GDPR monitoring purposes.

The Committee requests you to inform the REC once substantive changes to the protocol are made, important changes to the research team take place or researchers are added to the research team.

Moreover, the Committee requests you to send the final report of the research containing

The Netherlands
Fax +31 (0)20 568 8444

ABN AMRO 40 50 05 970
ABN AMRO USD 62 62 48 183

Royal Tropical Institute

Annex 6

Bar chart of Type of insurance among 450 respondents age > 60 years old in survey in Eastern Thailand 2018

